

23 April 2014

List of Marine Mammal Species and Subspecies

The [Committee on Taxonomy](#), chaired by Bill Perrin, has produced the first official SMM list of marine mammal species and subspecies. Consensus on some issues was not possible; this is reflected in the footnotes.

This list will be revisited and possibly revised every few months reflecting the continuing flux in marine mammal taxonomy. This version was updated on 23 April 2014.

This list can be cited as follows: "Committee on Taxonomy. 2014. List of marine mammal species and subspecies. Society for Marine Mammalogy, www.marinemammalscience.org, consulted on [date]."

This list includes living and recently extinct species and subspecies. It is meant to reflect prevailing usage and recent revisions published in the peer-reviewed literature. Author(s) and year of description of the species follow the Latin species name; when these are enclosed in parentheses, the species was originally described in a different genus. Classification and scientific names follow Rice (1998), with adjustments reflecting more recent literature. Common names are arbitrary and change with time and place; one or two currently frequently used in English and/or a range language are given here. Additional English common names and common names in French, Spanish, Russian and other languages are available at www.marinespecies.org/cetacea/.

Based on molecular and morphological data, the cetaceans genetically and morphologically fall firmly within the *artiodactyl clade* (Geisler and Uhen, 2005), and therefore we include them in the order *Cetartiodactyla*, with *Cetacea*, *Mysticeti* and *Odontoceti* as unranked taxa (recognizing that the classification within *Cetartiodactyla* remains partially unresolved -- e.g., see Spaulding et al., 2009, Price et al., 2005; Agnarsson and May-Collado, 2008)1. Below the rank of order, we list only families, species and subspecies, omitting superfamilies, subfamilies and taxa of other ranks.

For pinnipeds we previously followed Berta and Churchill (2012). To avoid issues of paraphyly, these authors proposed that based on data from genetics and morphology the genus *Arctocephalus* be limited to *Arctocephalus pusillus*, the type species of the genus *Arctocephalus*, and transferred the remaining 'Arctocephalus' species (i.e. *A. australis*, *A. galapagoensis*, *A. gazelle*, *A. philippii* and *A. tropicalis*) to *Arctophoca Peters*, 1866. However, Nyakatura and Bininda-Emonds (2012) compiled a new supertree of the *Carnivora* and concluded that this usage of *Arctophoca* may be premature because of remaining uncertainty about phylogenetic relationships, and we return provisionally to use of *Arctocephalus* for all the southern fur seals. Four subspecies of *A. australis* were formerly listed here: *A. a. australis*, *A. a. forsteri*, *A. a. gracilis* and *A. a. un-named*. However, Oliveira and Brownell (in press) synonymized *A. a. gracilis* with *A. a. australis*. The super-tree analysis by Nyakatura and Bininda-Emonds (2012) accords with the phylogenetic analysis of Higdon (2007), suggesting that the New Zealand fur seal should be recognized as a full species, *A. forsteri*. Two subspecies of *A. philippii* are valid: *A. p. philippii* and *A. p. townsendi*, although small sample sizes and a small number of genes sampled are concerns. Two subspecies of *Eumetopias* are supported largely on genetic data, which is also the case for recognition of California, Japanese and Galapagos sea lions as separate species. Brunner (2004) advised use of *Otaria byronia* (Blainville, 1820) over *O. flavescens* (Shaw, 1800). Lindqvist et al. (2009) concluded that a purported third subspecies of the walrus *Odobenus rosmarus laptevi* is not valid. Recent genetic analyses indicate that *Phoca vitulina concolor* is paraphyletic and this along with lack of morphological differentiation suggests that the western Atlantic subspecies is not valid; *P. v. vitulina* is considered here to apply to all Atlantic harbor seals. Within the North Pacific, until the subspecies limits of various populations are assessed only a single subspecies is recognized, *Phoca vitulina richardii*. Placement of the ringed seal, Caspian seal and Baikal seal has alternated between the genera *Phoca* and *Pusa*. We accept Rice's (1998) use of *Pusa* as the correct classification." The use of *Lontra* rather than *Lutra* for the marine otter follows Larivière (1998) in recognizing the otters of North and South America as a monophyletic taxon distinct from the otters of

Eurasia.

In the *mysticete* cetaceans, genetic evidence strongly supports the recognition of three separate phylogenetic species of right whales (Rosenbaum et al., 2000; Gaines et al., 2005). In addition, the genus *Eubalaena* (rather than *Balaena* as in Rice, 1998) is retained for the right whales as recommended by the Scientific Committee of the International Whaling Commission (IWC, 2002). *Caperea marginata* may be a member of the family *Cetotheriidae* (Fordyce and Marx 2012). *Neobalaenidae* is retained here provisionally. All Bryde's whales are provisionally considered to comprise a single species, *Balaenoptera edeni*, following the usage of the IWC (IWC 2002, 2008) and Kato and Perrin (2009). Some workers recognize *B. edeni* as including only the small-form coastal Bryde's whales of the western Pacific and Indian Oceans, using *B. brydei* for the globally distributed larger more oceanic form (Sasaki et al., 2006). Kato and Perrin (2009) and Kershaw et al. (2013) considered these more likely to be distinct at the subspecific level (although arguably at the species level), and they are included here provisionally as such. *Balaenoptera omurai* is a newly described species (Wada et al., 2003). It was previously confounded with the Bryde's whale and has been confirmed as having a separate and ancient lineage (Sasaki et al. 2006). Clarke (2004) proposed recognition of a pygmy form of the fin whale as a subspecies, based on distribution, size and coloration. He resurrected the synonym *patachonica* Burmeister, 1865 to apply to the subspecies: *B. physalus patachonica*. Branch et al. (2007) recognized the Chilean blue whale as an unnamed subspecies of *B. musculus*.

In the odontocetes, *Mesoplodon traversii* (spade-toothed whale) has been recognized as the senior synonym for *M. bahamondi* (Bahamonde's beaked whale) (van Helden et al., 2002). The first complete specimen was recently described from a stranding on the North Island of New Zealand (Thompson et al. 2012). *Mesoplodon perrini* is a newly described species (Dalebout et al., 2002). Dalebout et al. (in press) resurrected *Mesoplodon hotaula Deraniyagala*, 1963, a species closely similar to *M. ginkgodens*. The species *Inia boliviensis* d'Orbigny 1834 of the Cochabamba, Santa Cruz, Beni and Pando areas of the Bolivian Amazon basin is included in accordance with prevailing usage (Ruiz-García and Shostell, 2010). While the two *Inia* species overlap in all morphological characters (da Silva, 1994; Ruiz-García et al., 2006), they have been reproductively isolated from each other by a long series of rapids for an estimated 3.1 million years (Hollatz et al., 2011), and two independent lines of genetic evidence, from mtDNA and nuclear introns (Banguera-Hinestroza et al., 2008; Ruiz-García et al., 2008), suggest that they are on separate evolutionary trajectories and deserve recognition as phylogenetic species. Robineau et al. (2007) described the subspecies *Cephalorhynchus commersonii kerguelenensis*, and A. Baker et al. described *C. hectori maui*. The tucuxi has been split into the freshwater *Sotalia fluviatilis* (retaining the common name tucuxi) and the marine Guiana dolphin *S. guianensis* (Caballero et al. 2007). Based on a combined analysis of genetic and morphological data, Mendez et al. (2013) propose to recognize four species of Sousa (the humpback dolphins): the previously here-listed *S. teuszii* and *S. chinensis* plus *S. plumbea* and a new unnamed species from northern Australia originally documented based on molecular data by Frère et al. (2008). A drawback of the phylogenetic analyses is that there was only one sample from the area of supposed sympatry of *S. plumbea* and *S. chinensis* and very low coverage of the Indo-Malay region (n=5). The two species are listed here provisionally, pending the outcome of further analysis including more samples from those areas. The Burrnun dolphin *Tursiops australis*, recently described by Charlton-Robb et al. (2011), is not included here; its validity is uncertain. Among potential problems relating to its putative species status:

1. The specimens were compared morphologically only with bottlenose dolphins from Australia,
2. Despite the small sample sizes, the series overlapped in all metric characters and separation was possible only with multivariate analysis (which commonly resolves geographical forms within a species, e.g see Perrin et al. (1999) and Perrin et al. (2011) for *Stenella longirostris* and *Tursiops truncatus*, respectively),
3. Comparisons of external morphology and non-metrical characters were made only with *T. truncatus*, to the exclusion of *T. aduncus*,

4. Support for important nodes in molecular trees suggesting phylogenetic separation was low.

A rigorous re-evaluation of the relevant data and arguments is needed. Recognition of the Black Sea bottlenose dolphin is now well-supported by genetic data (Viaud-Martinez et al., 2008), as is the Black Sea common dolphin (Natoli et al., 2008). *Delphinus tropicalis* is now considered a subspecies of *D. capensis* (Jefferson and Van Waerebeek, 2002). *Lagenorhynchus* is widely considered an unnatural (polyphyletic) taxon containing morphologically convergent species (Cipriano 1997, LeDuc et al. 1999, McGowen 2011), and application of the genera *Sagmatias* (for *L. obscurus*, *obliquidens*, *australis* and *cruciger*) and *Leucopleurus* (for *L. acutus*) have been suggested as appropriate and used by some workers. However, there is continuing disagreement about whether *australis* and *cruciger* should be included in *Cephalorhynchus* (which would necessitate a new genus for *obliquidens* and *obscurus*, as *australis* is the type species for the genus *Sagmatias*) and about whether *albirostris* and *acutus* are sister species (which would obviate the need for *Leucopleurus*). We therefore provisionally retain all the species in *Lagenorhynchus*. Harlin-Cognato (2010) recognized *L. o. posidonia* (Peru/Chile). She also recognized *L. o. superciliosus* (Lesson and Garnot, 1826) for the New Zealand subspecies, but the species identity of the figure in Lesson and Garnot is in question, and we retain use of "un-named New Zealand subspecies." It has been noted repeatedly, most recently by Perrin et al. (2013), that the delphinine genera *Stenella* and *Tursiops* are paraphyletic and that at present there is no molecular or morphological basis for satisfactory resolution of phylogenetic relationships in the subfamily. A possible solution would be to return all the species in *Tursiops*, *Sousa*, and *Stenella* to *Delphinus*, the genus in which they were first described, and place *Lagenorhynchus hosei* there as well. However, considering that this would obscure the proven close relationship of the present *Delphinus* species and of the *Sousa* species, the status quo is maintained here provisionally, pending the outcome of more definitive morphological and molecular studies. Hopefully a more natural classification will emerge. Perrin et al. (1999) established the subspecies *Stenella longirostris roseiventris*. The Irrawaddy dolphin was recently split into *O. brevirostris* and *O. heinsohni*, the Australian snubfin dolphin (Beasley et al., 2005). Krahn et al. (2004) recognized two un-named species of killer whales, the resident and transient forms. Other forms of killer whales in the North Pacific, North Atlantic and Antarctic Ocean may warrant recognition as separate subspecies or even species, but the taxonomy has not yet been fully clarified or agreed (Morin et al. 2010; Foote et al. 2009, 2013). Wang et al. (2008) and Jefferson and Wang (2011) established *Neophocaena asiaorientalis* as a full species, with two subspecies. Viaud-Martinez et al. (2007) concluded based on morphological and genetic evidence that *Phocoena phocoena relicta* is a valid subspecies.

We list the baiji *Lipotes vexillifer* as "possibly extinct" in conformance with the IUCN Red List, although extinction seems a certainty.

In the *Sirenia*, subspecies of the dugong are not currently recognized (Domning, 1996). However, no in-depth study has been undertaken to address the issue of subspecies.

For review of species concepts, see Reeves et al. (2004), Orr and Coyne (2004), de Queiroz (2007) and Perrin (2009). Perrin et al. (2009) reviewed the cetacean subspecies, but that review has not yet appeared in the peer-reviewed literature and is therefore not considered here; the subspecies (including for the *Carnivora* and *Sirenia*) are as recognized by Rice (1998), with the above-noted changes.

Corrections and comments should be directed to the Committee on Taxonomy (william.perrin@noaa.gov). Divergent opinions by members of the Committee on particular taxonomic questions are given in the footnotes.

Order *Carnivora*

Family *Otariidae* (eared seals and sea lions; 15 species, of which 1 extinct)

Arctocephalus pusillus (Schreber, 1775) Cape fur seal

A. p. pusillus (Schreber, 1775). Cape fur seal

A. p. doriferus Wood Jones, 1925. Australian fur seal

Arctocephalus gazella (Peters, 1876). Antarctic fur seal

Arctocephalus tropicalis (Gray, 1872). Subantarctic fur seal

Arctocephalus forsteri (Lesson, 1828). New Zealand fur seal

Arctocephalus australis (Zimmermann, 1783) South American fur seal

A. a. australis (Zimmermann, 1783). South American fur seal

A. a. un-named subspecies. Peruvian fur seal

Arctocephalus galapagoensis Heller, 1904. Galapagos fur seal

Arctocephalus philippii (Peters, 1866)

A. p. philippii Peters, 1866. Juan Fernandez fur seal

A. p. townsendi (Merriam, 1897). Guadalupe fur seal

Callorhinus ursinus (Linnaeus, 1758). Northern fur seal

Zalophus japonicus (Peters, 1866). Japanese sea lion (extinct)

Zalophus californianus (Lesson, 1828). California sea lion

Zalophus wollebaeki Sivertsen, 1953. Galapagos sea lion

Eumetopias jubatus (Schreber, 1776). Steller sea lion, northern sea lion

E. j. jubatus (Schreber, 1776). Western Steller sea lion

E. j. monteriensis (Gray, 1859). Loughlin's Steller sea lion

Neophoca cinerea (Peron, 1816). Australian sea lion

Phocarctos hookeri (Gray, 1844). New Zealand sea lion

Otaria byronia (Blainville, 1820). South American sea lion

Family *Odobenidae*

Odobenus rosmarus (Linnaeus, 1758). Walrus

O. r. rosmarus (Linnaeus, 1758). Atlantic walrus

O. r. divergens (Illiger, 1815). Pacific walrus

Family *Phocidae* (earless seals; 19 species, of which 1 extinct)

Erignathus barbatus (Erxleben, 1777). Bearded seal

E. b. barbatus (Erxleben, 1777). Atlantic bearded seal

E. b. nauticus (Pallas, 1881). Pacific bearded seal

Phoca vitulina (Linnaeus, 1758). Harbor seal, common seal

P. v. vitulina (Linnaeus, 1758). Atlantic harbor seal

P. v. mellonae (Doutt, 1942). Ungava harbor seal

P. v. richardii (Gray, 1864). Pacific harbor seal

Phoca largha (Pallas, 1811). Spotted seal, largha seal

Pusa hispida (Schreber, 1775). Ringed seal

P. h. hispida (Schreber, 1775). Arctic Ringed seal

P. h. botnica (Gmelin, 1788). Baltic ringed seal

P. h. ochotensis (Nordquist, 1889) Okhotsk ringed seal

P. h. ladogensis (Nordquist, 1889). Lake Ladoga seal

P. h. saimensis (Nordquist, 1889). Saima seal

Pusa caspica (Gmelin, 1788). Caspian seal

Pusa sibirica (Gmelin, 1788). Baikal seal

Halichoerus grypus (Fabricius, 1791). Gray seal

H. g. grypus (Fabricius, 1791). Western Atlantic gray seal

H. g. macrorhynchus Hornschuh and Schilling, 1851. Eastern Atlantic gray seal

Histiophoca fasciata (Zimmerman, 1783). Ribbon seal

Pagophilus groenlandicus (Erxleben, 1777). Harp seal

Cystophora cristata (Erxleben, 1777). Hooded seal

Monachus tropicalis (Gray, 1850). Caribbean monk seal (extinct)

Monachus monachus (Hermann, 1779). Mediterranean monk seal

Monachus schauinslandi Matschie, 1905. Hawaiian monk seal

Mirounga leonina (Linnaeus, 1758). Southern elephant seal

Mirounga angustirostris (Gill, 1866). Northern elephant seal

Leptonychotes weddellii (Lesson, 1826). Weddell seal

Ommatophoca rossii Gray, 1844. Ross seal

Lobodon carcinophaga (Hombron and Jacquinot, 1842). Crabeater seal

Hydrurga leptonyx (Blainville, 1820). Leopard seal

Family *Ursidae*

Ursus maritimus Phipps, 1774. Polar bear

U. m. maritimus Phipps, 1774. Atlantic polar bear

U. m. marinus Pallas, 1776. Pacific polar bear

Family *Mustelidae*

Enhydra lutris (Linnaeus, 1758). Sea otter

E. l. lutris (Linnaeus, 1758). Western sea otter

E. l. kenyonii Wilson, 1991. Eastern sea otter

E. l. nereis (Merriam, 1904). Southern sea otter

Lontra felina (Molina, 1782). Chungungo, marine otter

Neovison macrodon (Prentis, 1903). Sea mink (extinct)

Order *Cetartiodactyla* (artiodactyls and cetaceans)

CETACEA (cetaceans; 90 species, of which 1 possibly extinct)

MYSTICETI (baleen whales, 14 species)

Family *Balaenidae* (right whales, 4 species)

Eubalaena glacialis (Müller, 1776). North Atlantic right whale

Eubalaena japonica (Lacépède, 1818). North Pacific right whale

Eubalaena australis (Desmoulins, 1822). Southern right whale

Balaena mysticetus Linnaeus, 1758. Bowhead whale, Greenland whale

Family *Neobalaenidae*

Caperea marginata (Gray, 1846). Pygmy right whale

Family *Eschrichtiidae*

Eschrichtius robustus (Lilljeborg, 1861). Gray whale

Family *Balaenopteridae* (rorquals, 8 species)

Megaptera novaeangliae (Borowski, 1781). Humpback whale

Balaenoptera acutorostrata Lacépède, 1804. Common minke whale

B. a. acutorostrata Lacépède, 1804. North Atlantic minke whale

B. a. scammoni Deméré, 1986. North Pacific minke whale

B. a. un-named subsp. Dwarf minke whale

Balaenoptera bonaerensis Burmeister, 1867. Antarctic minke whale

Balaenoptera edeni Anderson, 1879. Bryde's whale

B. e. edeni Anderson, 1879. Eden's whale

B. e. brydei Olsen, 1913. Offshore Bryde's whale

Balaenoptera omurai Wada, Oishi and Yamada, 2003. Omura's whale

Balaenoptera borealis Lesson, 1828. Sei whale

B. b. borealis Lesson, 1828. Northern sei whale

B. b. schlegellii (Flower, 1865). Southern sei whale

Balaenoptera physalus (Linnaeus, 1758). Fin whale

B. p. physalus (Linnaeus, 1758). Northern fin whale

B. p. quoyi Fischer, 1829). Southern fin whale

B. p. patachonica Burmeister, 1865. Pygmy fin whale

Balaenoptera musculus (Linnaeus, 1758). Blue whale

B. m. musculus (Linnaeus, 1758). Northern blue whale

B. m. intermedia Burmeister, 1871. Antarctic blue whale

B. m. indica Blyth, 1859. Northern Indian Ocean blue whale

B. m. brevicauda Ichihara, 1966. Pygmy blue whale

B. m. un-named subsp. Chilean blue whale.

ODONTOCETI (toothed whales, dolphins and porpoises: 76 named and 1 un-named species; one named species possibly extinct)

Family *Physeteridae*

Physeter macrocephalus Linnaeus, 1758. Sperm whale, cachalot

Family *Kogiidae*

Kogia breviceps (Blainville, 1838). Pygmy sperm whale

Kogia sima (Owen, 1866). Dwarf sperm whale

Family *Ziphiidae* (beaked whales, 22 species)

Ziphius cavirostris G. Cuvier, 1823. Cuvier's beaked whale, goose-beaked whale

Berardius arnuxii Duvernoy, 1851. Arnoux' beaked whale

Berardius bairdii Stejneger, 1883. Baird's beaked whale

Tasmacetus shepherdi Oliver, 1937. Shepherd's beaked whale, Tasman beaked whale

Indopacetus pacificus (Longman, 1926). Longman's beaked whale, tropical bottlenose whale,

Hyperoodon ampullatus (Forster, 1770). Northern bottlenose whale

Hyperoodon planifrons Flower, 1882. Southern bottlenose whale

Mesoplodon hectori (Gray, 1871). Hector's beaked whale

Mesoplodon mirus True, 1913. True's beaked whale

Mesoplodon europaeus (Gervais, 1855). Gervais' beaked whale

Mesoplodon bidens (Sowerby, 1804). Sowerby's beaked whale

Mesoplodon grayi von Haast, 1876. Gray's beaked whale

Mesoplodon perrini Dalebout, Mead, Baker, Baker and van Helden, 2002. Perrin's beaked whale

Mesoplodon peruvianus Reyes, Mead and Van Waerebeek, 1991. Pygmy beaked whale

Mesoplodon bowdoini Andrews, 1908. Andrews' beaked whale

Mesoplodon traversii (Gray, 1874). Spade-toothed whale

Mesoplodon carlhubbsi Moore, 1963. Hubbs' beaked whale

Mesoplodon ginkgodens Nishiwaki and Kamiya, 1958. Ginkgo-toothed beaked whale

Mesoplodon stejnegeri True, 1885. Stejneger's beaked whale

Mesoplodon layardii (Gray, 1865). Strap-toothed beaked whale, Layard's beaked whale

Mesoplodon densirostris (Blainville, 1817. Blainville's beaked whale

Mesoplodon hotaula Deraniyagala, 1963. Deraniyagala's beaked whale.

Family *Platanistidae*

Platanista gangetica (Lebeck, 1801). South Asian river dolphin, Indian river dolphin

P. g. gangetica (Lebeck, 1801). Susu, Ganges river dolphin

P. g. minor Owen, 1853. Bhulan, Indus river dolphin

Family *Iniidae*

Inia geoffrensis (Blainville, 1817). Amazon river dolphin

I. g. geoffrensis (Blainville, 1817). Boto

I. g. humboldtiana Pilleri and Gühr, 1977. Orinoco bufeo

Inia boliviensis d'Orbigny, 1834. Bolivian bufeo

Family *Lipotidae*

Lipotes vexillifer (Miller, 1918). Baiji, Yangtze river dolphin – possibly extinct

Family *Pontoporiidae*

Pontoporia blainvillei (Gervais and d'Orbigny, 1844). Franciscana, toninha.

Family *Monodontidae*

Monodon monoceros Linnaeus, 1758. Narwhal

Delphinapterus leucas (Pallas, 1776). Beluga, white whale

Family *Delphinidae* (38 species)

Cephalorhynchus commersonii (Lacépède, 1804). Commerson's dolphin

C. c. commersonii (Lacépède, 1804). Commerson's dolphin

C. c. kerguelenensis Robineau, Goodall, Pichler and C. S. Baker, 2007. Kerguelen Islands Commerson's dolphin

Cephalorhynchus eutropia (Gray, 1846). Chilean dolphin

Cephalorhynchus heavisidii (Gray, 1828). Heaviside's dolphin, Haviside's dolphin

Cephalorhynchus hectori (Van Beneden, 1881). Hector's dolphin, New Zealand dolphin

C. h. hectori (Van Beneden, 1881). South Island Hector's dolphin

C. h. maui A. Baker, Smith and Pichler, 2002. Maui's dolphin, North Island Hector's dolphin

Steno bredanensis (G. Cuvier in Lesson, 1828). Rough-toothed dolphin

Sousa teuszii (Kükenthal, 1892). Atlantic humpback dolphin

Sousa chinensis (Osbeck, 1765). Pacific humpback dolphin

Sousa plumbea (G. Cuvier, 1829). Indian Ocean humpback dolphin

Sousa un-named species. Australian humpback dolphin

Sotalia fluviatilis (Gervais and Deville in: Gervais, 1853). Tucuxi

Sotalia guianensis (Van Bénédén, 1864). Guiana dolphin, costero

Tursiops truncatus (Montagu, 1821). Common bottlenose dolphin

T. t. truncatus (Montagu, 1821). Common bottlenose dolphin

T. t. ponticus Barabash-Nikiforov, 1940. Black Sea bottlenose dolphin

Tursiops aduncus (Ehrenberg, 1833). Indo-Pacific bottlenose dolphin

Stenella attenuata (Gray, 1846). Pantropical spotted dolphin

S. a. attenuata (Gray, 1846). Offshore pantropical spotted dolphin

S. a. graffmani (Lönnerberg, 1934). Coastal pantropical spotted dolphin

Stenella frontalis (G. Cuvier, 1829). Atlantic spotted dolphin

Stenella longirostris (Gray, 1828). Spinner dolphin

S. l. longirostris (Gray, 1828). Gray's spinner dolphin

S. l. orientalis Perrin, 1990. Eastern spinner dolphin

S. l. centroamericana Perrin, 1990. Central American spinner dolphin

S. l. roseiventris (Wagner, 1846). Dwarf spinner dolphin

Stenella clymene (Gray, 1850). Clymene dolphin

Stenella coeruleoalba (Meyen, 1833). Striped dolphin

Delphinus delphis Linnaeus, 1758. Short-beaked common dolphin, saddleback dolphin

D. d. delphis Linnaeus, 1758. Short-beaked common dolphin

D. d. ponticus Barabash, 1935. Black Sea common dolphin

Delphinus capensis Gray, 1828. Long-beaked common dolphin

D. c. capensis Gray, 1828. Long-beaked common dolphin

D. c. tropicalis van Bree, 1971. Indo-Pacific common dolphin

Lagenodelphis hosei Fraser, 1956. Fraser's dolphin

Lagenorhynchus albirostris (Gray, 1846). White-beaked dolphin

Lagenorhynchus acutus (Gray, 1828). Atlantic white-sided dolphin

Lagenorhynchus obliquidens Gill, 1865. Pacific white-sided dolphin

Lagenorhynchus obscurus (Gray, 1828). Dusky dolphin

L. o. obscurus (Gray, 1828). African dusky dolphin

L. o. fitzroyi (Waterhouse, 1838). Fitzroy's dolphin

L. o. posidonia (Philippi, 1893). Peruvian/Chilean dusky dolphin

L. o. un-named subsp. New Zealand dusky dolphin

Lagenorhynchus australis (Peale, 1848). Peale's dolphin

Lagenorhynchus cruciger (Quoy and Gaimard, 1824). Hourglass dolphin

Lissodelphis borealis (Peale, 1848). Northern right-whale dolphin

Lissodelphis peronii (Lacépède, 1804). Southern right-whale dolphin

Grampus griseus (G. Cuvier, 1812). Risso's dolphin, gray grampus

Peponocephala electra (Gray, 1846). Melon-headed whale, Electra dolphin

Feresa attenuata Gray, 1874. Pygmy killer whale

Pseudorca crassidens (Owen, 1846). False killer whale

Orcinus orca (Linnaeus, 1758). Killer whale, orca

O. o. un-named subsp. Resident killer whale

O. o. un-named subsp. Transient killer whale, Bigg's killer whale

Globicephala melas (Traill, 1809). Long-finned pilot whale

G. m. melas (Traill, 1809). North Atlantic long-finned pilot whale

G. m. edwardii (A. Smith, 1834). Southern long-finned pilot whale

G. m. un-named subsp. North Pacific long-finned pilot whale

Globicephala macrorhynchus Gray, 1846. Short-finned pilot whale

Orcaella brevirostris (Owen in Gray, 1866). Irrawaddy dolphin, pesut

Orcaella heinsohni Beasley, Robertson and Arnold, 2005. Australian snubfin dolphin

Family *Phocoenidae* (porpoises, 7 species)

Neophocaena phocaenoides (G. Cuvier, 1829). Indo-Pacific finless porpoise

Neophocaena asiaorientalis (Pilleri and Gühr, 1972). Narrow-ridged finless porpoise

N. a. asiaorientalis (Pilleri and Gühr, 1972). Yangtze finless porpoise

N. a. sunameri Pilleri and Gühr, 1975. East Asian finless porpoise, sunameri

Phocoena phocoena (Linnaeus, 1758). Harbor porpoise, common porpoise

P. p. phocoena (Linnaeus, 1758). Atlantic harbor porpoise

P. p. vomerina (Gill, 1865). Eastern Pacific harbor porpoise

P. p. relicta Abel, 1905. Black Sea harbor porpoise

P. p. un-named subsp. Western Pacific harbor porpoise

Phocoena sinus Norris and McFarland, 1958. Vaquita, Gulf of California harbor porpoise

Phocoena spinipinnis Burmeister, 1865. Burmeister's porpoise

Phocoena dioptrica Lahille, 1912. Spectacled porpoise

Phocoenoides dalli (True, 1885). Dall's porpoise, Dall porpoise

P. d. dalli (True, 1885). *Dalli*-type Dall's porpoise

P. d. truei Andrews, 1911. *Truei*-type Dall's porpoise

ORDER SIRENIA (sirenians, 5 species – 1 extinct)

Family Trichechidae

Trichechus manatus Linnaeus, 1758. West Indian manatee

T. m. manatus Linnaeus, 1758. Antillean manatee

T. m. latirostris (Harlan, 1824). Florida manatee

Trichechus senegalensis Link, 1795. West African manatee, African manatee

Trichechus inunguis (Natterer, 1883). Amazonian manatee

Family Dugongidae

Dugong dugon (Müller, 1776). Dugong

Hydrodamalis gigas (Zimmerman, 1780). Steller's sea cow - extinct

Footnotes

¹Use of Order *Cetartiodactyla* is favored by most evolutionary mammalogists working with molecular data. Some others, including many marine mammalogists and paleontologists, favor retention of Order *Cetacea* in the interest of taxonomic stability.

²(from D. Rice) Baker *et al.* (2003) hold that there is no evidence that would support the classification of

the right whales as more than a single biological species. [The three species are here recognized as phylogenetic species.]

References

Agnarsson, I. and L. J. May-Collado. 2008. The phylogeny of *Cetartiodactyla*: the importance of dense taxon sampling, missing data, and the remarkable promise of cytochrome b to provide reliable species-level phylogenies. *Molecular Phylogenetics and Evolution* 48:964–985.

Baker, A. N., A. N. H. Smith, and F. B. Pichler. 2002. Geographical variation in Hector's dolphin : recognition of a new subspecies of *Cephalorhynchus hectori*. *Journal of the Royal Society of New Zealand* 32:713–727.

Baker, R. J., L. C. Bradley, R. D. Bradley, J. W. Dragoo, M. D., R. S. Hoffman, C. A. Jones, F. Reid, D. W. Rice and C. Jones. 2003. Revised checklist of North American mammals north of Mexico, 2003. *Museum of Texas Tech University Occasional Papers* 229:1–24.

Banguera-Hinestroza, E., M. Cárdenas, M. Ruiz-García, M. Marmontel, E. Gaitán, R. Vázquez and F. García-Vallejo. 2008. Molecular identification of evolutionarily significant units in the Amazon river dolphin *Inia* sp. (Cetacea: Iniidae). *Journal of Heredity* 93:313–322.

Beasley, I., K. M. Robertson and P. Arnold. 2005. Description of a new dolphin, the Australian snubfin dolphin *Orcaella heinsohni* sp. n. (Cetacea: Delphinidae). *Marine Mammal Science* 21: 365–400.

Berta, A. and M. Churchill. 2012. Pinniped taxonomy: review of currently recognized species and subspecies, and evidence used for their description. *Mammal Review* 42:207–234.

Branch, T. A., E. M. N. Abubaker, S. Mkango and D. S. Butterworth. 2007. Separating southern blue whale subspecies based on length frequencies of sexually mature females. *Marine Mammal Science* 23:803–833.

Brunner, S. 2004. Fur seals and sea lions (*Otariidae*): identification of species and taxonomic review. *Systematics and Biodiversity* 1:339–439.

Caballero, S., F. Trujillo, J. A. Vianna, H. Barrios-Garrido, M. G. Montiel, S. Beltrán-Pedrerros, M. Marmontel, M. C. Santos, M. Rossi-Santos, F. R. Santos, and C. S. Baker. 2007. Taxonomic status of the genus *Sotalia*: species-level ranking for "tucuxi" (*Sotalia fluviatilis*) and "costero" (*Sotalia guianensis*) dolphins. *Marine Mammal Science* 23: 358–386.

Charlton-Robb, K., L. Gershwin, R. Thompson, J. Austin, K. Owen and S. McKechnie. 2011. A new dolphin species, the Burrnunan dolphin *Tursiops australis* sp. Nov., endemic to southern Australian waters. *PLoS ONE* 6(0): e24047.doi:10.1371/journal.pone.0024047.

Cipriano, F. 1997. Antitropical distributions and speciation in dolphins of the genus *Lagenorhynchus*: a preliminary analysis. Pages 305–316 in A. E. Dizon, S. J. Chivers and W. F. Perrin (eds). *Molecular genetics of marine mammals*. Society for Marine Mammalogy Special Publication 3.

Clarke, R. 2004. Pygmy fin whales. *Marine Mammal Science* 20:329–334.

Dalebout, M. L., J. G. Mead, C. S. Baker, A. N. Baker, and A. L. van Helden. 2002. A new species of beaked whale *Mesoplodon perrini* sp. n. (Cetacea: Ziphiidae) discovered through phylogenetic analyses of mitochondrial DNA sequences. *Marine Mammal Science* 18:577–608.

- Dalebout, M. L., S. C. Baker, D. Steel, K. Thompson, K. M. Robertson, S. J. Chivers, W. F. Perrin, M. Goonatilake, R. C. Anderson, J. G. Mead, C. W. Potter, L. Thompson, D. Jupiter and T. K. Yamada. In press. Resurrection of *Mesoplodon hotaula* Deraniyagala 1963: a new species of beaked whale in the tropical Indo-Pacific. *Marine Mammal Science*.
- Dasmahapatra, K. K., J. I. Hoffman and W. Amos. 2009. Pinniped phylogenetic relationships inferred using AFLP markers. *Heredity* 103:168—177.
- de Queiroz, K. 2007. Species concepts and species delineation. *Systematic Biology* 56:879—886.
- Domning, D. 1996. Bibliography and index of the Sirenia and Desmostylia. *Smithsonian Contributions to Paleobiology* 80:1—611.
- Foot, A. D., J. Newton, S. B. Piertney, E. Willersley, and M. T. P. Gilbert. 2009. Ecological, morphological and genetic divergence of sympatric North Atlantic killer whale populations. *Molecular Ecology* 18:5207—5217.
- Foot, A. D., J. Newton, M. C. Ávila-Arcos, M.-L. Kampmann, J. A. Samaniego, K. Post, A. Rosing-Asvid, M.-H.S. Sinding and M. T. P. Gilbert. 2013. Tracking niche variation over millennial timescales in sympatric killer whale lineages. *Proceedings of the Royal Society B*. 280:20131481.
- Fordyce, R. E. and F. G. Marx. 2012. The pygmy right whale *Caperea marginata*: last of the cetotheres. *Proceedings of the Royal Society B* 280:20122645.
- Frère, C. H. P. T. Porter, V. G. Cockcroft and M. L. Dalebout. 2008. Phylogenetic analysis of mtDNA sequences suggests revision of humpback dolphin (*Sousa* spp.) taxonomy is needed. *Marine and Freshwater Research* 59:259—268.
- Gaines, C. A., M. P. Hare, S. E. Beck and H. C. Rosenbaum. 2005. Nuclear markers confirm taxonomic status and relationships among highly endangered and closely related right whale species. *Proceedings of the Royal Society B* 272:533—542.
- Geisler, J. H. and M. D. Uhen. 2005. Phylogenetic relationships of extinct cetartiodactyls: results of simultaneous analyses of molecular, morphological, and stratigraphic data. *Journal of Mammalian Evolution* 12:145--160.
- Harlin-Cognato, A. D. 2010. The dusky dolphin's place in the delphinid family tree. Pages 1—20 *in* B. Würsig and M. Würsig (eds). *The dusky dolphin. Master acrobat off different shores*. Academic Press, Amsterdam.
- van Helden, A. L., A. N. Baker, M. L. Dalebout, J. C. Reyes, K. Van Waerebeek, and C. S. Baker. 2002. Resurrection of *Mesoplodon traversii* (Gray, 1874), senior synonym of *M. bahamondi* Reyes, Van Waerebeek, Cárdenas and Yáñez, 1995 (Cetacea: Ziphiidae). *Marine Mammal Science* 18:609-621.
- Higdon, J. W., O. R. P. Bininda-Emonds, R. M. Beck and S. H. Ferguson. 2007. Phylogeny and divergence of the pinnipeds (Carnivora, Mammalia) assessed using a multigene dataset. *BMC Evolutionary Biology* 7:216.
- Hollatz, C., S. Torres Vilaça, R. A. F. Redondo, M. Marmontel, C. S. Baker and F. R. Santos. 2011. The Amazon River system as an ecological barrier driving genetic differentiation of the pink dolphin (*Inia*

geoffrensis). Biological Journal of the Linnaen Society 102:812—827.

International Whaling Commission. 2001. Report of the Scientific Committee. Journal of Cetacean Research and Management 3, Supplement:1--75.

International Whaling Commission. 2008. Report of the Scientific Committee. Journal of Cetacean Research and Management 10, Supplement: 1--406.

Jefferson, T. A. and K. Van Waerebeek. 2002. The taxonomic status of the nominal species *Delphinus tropicalis* van Bree, 1971. Marine Mammal Science 18:787--818.

Jefferson, T. A. and J. Y. Wang. 2011. Revision of the taxonomy of finless porpoises (genus *Neophocaena*): the existence of two species. Journal of Marine Animals and Their Ecology 4:3—16.

Kato, H. and W. F. Perrin. 2009. Bryde's whales *Balaenoptera edeni/brydei*. Pages 158—163 in W. F. Perrin, B. Würsig and J. G. M. Theewissen, eds. Encyclopedia of Marine Mammals. Academic Press, Amsterdam.

Kershaw, F., M. S. Leslie, T. Collins, R. M. Mansur, B. D. Smith, G. Minton, R. Baldwin, R. G. LeDuc, R. C. Anderson, R. L. Brownell, Jr. and H. C. Rosenbaum. 2013. Population differentiation of 2 forms of Bryde's whales in the Indian and Pacific Oceans. Journal of Heredity doi:10.1093/jhered/est057. 10pp.

Krahn, M., M. J. Ford, W. F. Perrin, P. R. Wade, R. P. Angliss, M. B. Hanson, B. L. Taylor, G. M. Ylitalo, M. E. Dahlheim, J. E. Stein and R. S. Waples. 2004 Status Review of Southern Resident Killer Whales (*Orcinus orca*) under the Endangered Species Act. NOAA Technical Memorandum NMFS-NWFSC-62. 73pp.

Larivière, S. 1998. *Lontra felina*. Mammalian Species 575:1--5.

Lindqvist, C., L. Bachmann, L. W. Andersen, E. W. Born, U. Arnason, K. M. Kovacs, C. Lydersen, A. V. Abramov and Ø. Wiig. 2008. The Laptev Sea walrus *Odobenus rosmarus laptevi*: an enigma revisited. Zoologica Scripta 38:113—127.

McGowen, M. R. 2011. Toward the resolution of an explosive radiation—A multilocus phylogeny of oceanic dolphins (Delphinidae). Molecular Phylogenetics and Evolution 60:345—357.

Mendez, M., T. A. Jefferson, S.-O. Kolokotronis, M. Krützen, G. J. Parra, T. Collins, G. Minton, R. Baldwin, P. Berggren, A. Särnblad, O. A. Amir, V. M. Peddemors, L. Karczmarski, A. Guissamulo, B. Smith, D. Sutaria, G. Amato and H. C. Rosenbaum. 2013. Integrating multiple lines of evidence to better understand the evolutionary divergence of humpback dolphins along their entire distribution range: a new dolphin species in Australian waters. Molecular Ecology doi: 10.1111/mec.12535.

Morin, P. A., F. I. Archer, A. D. Foote, J. Vilstrup, E. A. Allen, P. Wade, J. Durban, K. Parsons, R. Pitman, L. Li, P. Bouffard, S. C. Abel Nielsen, M. Rasmussen, E. Willerslev, M. T. P. Gilbert and T. Harkins. 2010. Complete mitochondrial phylogenetic analysis of killer whales (*Orcinus orca*) indicates multiple species. Marine Genomic Research 20:908—916.

Natoli, A., A. Cañadas, C. Vaquero, E. Politi, P. Fernandez-Navarro and A. R. Hoelzel. 2008. Conservation genetics of the short-beaked common dolphin (*Delphinus delphis*) in the Mediterranean Sea and the eastern North Atlantic. Conservation Genetics 9:1479—1487.

Nyakatura, K. and O. R.P. Birinda-Emonds. 2013. Updating the evolutionary history of Carnivora

(Mammalia): a new species-level supertree complete with divergence time estimates. *BMC Biology* 10:1—31.

Oliveira, L. R. and R. L. Brownell, Jr. In press. Taxonomic status of two subspecies of South American fur seals: *Arctocephalus australis australis* vs. *A. a. gracilis*. *Marine Mammal Science*.

Orr, H. A. and J. A. Coyne. 2004. *Speciation*. Sinauer Associates, Sunderland, Massachusetts.

Parra, G. J. and G. J. B. Ross. 2009. Humpback dolphins *S. chinensis* and *S. teuszii*. Pages 576—582 in W. F. Perrin, B. Würsig and J. G. M. Thewissen, eds. *Encyclopedia of Marine Mammals*. Academic Press, Amsterdam.

Perrin, W. F. 2009. Species. Pages 1084—1087 in W. F. Perrin, B. Würsig and J. G. M. Thewissen, eds. *Encyclopedia of Marine Mammals*. Academic Press, Amsterdam.

Perrin, W. F., M. L. L. Dolar and D. Robineau. 1999. Spinner dolphins (*Stenella longirostris*) of the western Pacific and Southeast Asia: pelagic and shallow-water forms. *Marine Mammal Science* 15:1029—1053.

Perrin, W. F., J. G. Mead and R. L. Brownell, Jr. 2009. Review of the evidence used in the description of currently recognized cetacean subspecies. NOAA Technical Memorandum NOAA-TM-NMFS-SWFSC. 450. 35 pp..

Perrin, W. F., J. L. Thieleking, W. A. Walker, F. I. Archer and K. M. Robertson. 2011. Common bottlenose dolphins (*Tursiops truncatus*) in California waters: Cranial differentiation of coastal and offshore ecotypes. *Marine Mammal Science* 27:769—792,

Perrin, W. F., P. E. Rosel and F. Cipriano. 2013. How to contend with paraphyly in the taxonomy of the delphinine cetaceans. *Marine Mammal Science* 29:567—588.

Phillips, C. D., J. W. Bickham, J. C. Patton and T. S. Gelatt. 2009. Systematics of Steller sea lions (*Eumetopias jubatus*): subspecies recognition based on concordance of genetics and morphometrics. *Museum of Texas Tech University Occasional Papers* 283:1—15.

Price, S. A., O. R. P. Bininda-Emonds and J. L. Gittleman. 2005. A complete phylogeny of the whales, dolphins and even-toed hoofed mammals (Cetartiodactyla). *Biological Review* 80:445—473.

Reeves, R. R., W. F. Perrin, B. L. Taylor, C. S. Baker and S. L. Mesnick, eds. 2004. Report of the Workshop on Shortcomings of Cetacean Taxonomy in Relation to Needs of Conservation and Management, April 30—May 2, 2004 La Jolla, California. NOAA Technical Memorandum NOAA-TM-NMFS-SWFSC-363:1—94.

Rice, D. W. 1998. *Marine mammals of the world. Systematics and distribution*. Society for Marine Mammalogy Special Publication 4:1--231.

Robineau, D., R. N. P. Goodall, F. Pichler, and C. S. Baker. 2007. Description of a new subspecies of Commerson's dolphin, *Cephalorhynchus commersonii* (Lacépède, 1804) inhabiting the coastal waters of the Kerguelen Islands. *Mammalia* 2007:172—180.

Rosenbaum, H., R. L. Brownell, Jr., M. W. Brown, C. Schaeff, Y. Portway, B. N. White, S. Malik, L. A. Pastene, N. J. Patenaude, C. S. Baker, M. Goto, P. B. Best, P. J. Clapham, P. Hamilton, R. Payne, V. Rowntree, C. T. Tynan, J. L. Bannister, R. and DeSalle. 2000. World-wide genetic differentiation of

Eubalaena questioning the number of right whale species. *Molecular Ecology* 9:1793--1802.

Ruiz-García, M and J. M. Shostell (eds.). 2010. *Biology, evolution and conservation of river dolphins within South America and Asia*. Nova Science Publishers, New York.

Ruiz-García, M., E. Banguera and H. Cárdenas. 2006. Morphological analysis of three *Inia* (Cetacea: Iniidae) populations from Colombia and Bolivia. *Acta Theriologica* 51:411—426.

Ruiz-García, M., S. Caballero, M. Martínez-Agüero and J. M. Shostell. 2008. Molecular differentiation among *Inia geoffrensis* and *Inia boliviensis* (Iniidae, Cetacea) by means of nuclear intron sequences. Pages 1—25 in V. T. Koven, ed., *Population Genetics Research Progress Chapter 6*.

Sasaki, T., M. Nikaido, S. Wada, T. K. Yamada, Y. Cao, M. Hasegawa, and N. Okada. 2006. *Balaenoptera omurai* is a newly discovered baleen whale that represents an ancient evolutionary lineage. *Molecular Phylogenetics and Evolution* 41:40--52.

da Silva, V. M. F. 1994. *Aspects of the biology of the Amazonian dolphins genera Inia and Sotalia fluviatilis*. Ph.D. thesis, University of Cambridge, Cambridge, U.K. 328 pp.

Spaulding, M., M. A. O'Leary and J. Gatesy. 2009. Relationships of Cetacea (Artiodactyla) among mammals: increased taxon sampling alters interpretations of key fossils and character evolution. *PLoS ONE* 4:1—14.

Thompson, K., C. S. Baker, A. van Helden, S. Paatel, C. Millar and R. Constantine. 2012. The world's rarest whale. *Current Biology* 22:R905—R906.

Viaud-Martinez, M. Martínez Vergara, P. E. Gol'din, V. Ridoux, A. A., Öztürk, B. Öztürk, P. E. Rosel, A. Frantzis, A. Komnenou, and A. J. Bohanak. 2007. Morphological and genetic differentiation of the Black Sea harbour porpoise *Phocoena phocoena*. *Marine Ecology Progress Series* 338:281—294.

Viaud-Martinez, K. A., R. L. Brownell, Jr., A. Komnenou, and A. J. Bohanak. 2008. Genetic isolation and morphological divergence of Black Sea bottlenose dolphins. *Biological Conservation* 141:1600—1611.

Wada, S., M. Oishi and T. K. Yamada. 2003. A newly discovered species of living baleen whale. *Nature* 426:278--281.

Wang, J. Y., T. R. Frasier, S. C. Yang and B. N. White. 2008. Detecting recent speciation events: the case of the finless porpoise (genus *Neophocaena*). *Heredity* 101:145—155.



Last updated April 23 2014 by members of the [Committee on Taxonomy](#):

- William F. Perrin (Chair)
- C. Scott Baker
- Annalisa Berta
- Daryl J. Boness
- Robert L. Brownell, Jr.
- Daryl P. Domning
- R. Ewan Fordyce
- Rebecca M. Hamner

- Thomas A. Jefferson
- James G. Mead
- Larissa R. Oliveira
- Dale W. Rice
- Patricia E. Rosel
- John Y. Wang
- Tadasu Yamada