

# List of Marine Mammal Species and Subspecies

## Introduction

The [Committee on Taxonomy](#), chaired by Patricia Rosel, produced the first official Society for Marine Mammalogy list of marine mammal species and subspecies in 2010. Consensus on some issues has not been possible; this is reflected in the footnotes. The Terms of Reference provide information on the Committee's process. The list is updated at least annually.

The current version was updated in June 2023. This list can be cited as follows: "Committee on Taxonomy. 2023. List of marine mammal species and subspecies. Society for Marine Mammalogy, [www.marinemammalscience.org](http://www.marinemammalscience.org), consulted on [date]."

This [list](#) includes living and recently extinct (within historical times) species and subspecies. It is meant to reflect prevailing usage and recent revisions published in the peer-reviewed literature. Classification and scientific names follow Rice (1998), with adjustments reflecting more recent literature. Author(s) and year of description of each taxon follow the Latin (scientific) species name; when these are enclosed in parentheses, the taxon was originally described in a different genus.

The Committee considers and evaluates new, peer-reviewed literature that proposes taxonomic changes. The Committee's focus is on alpha taxonomy (describing and naming taxa) and beta taxonomy primarily at lower levels of the hierarchy (subspecies, species and genera), although it may evaluate issues at higher levels if deemed necessary. Proposals for new, taxonomically distinct taxa require a formal, peer-reviewed study and should provide robust evidence that some subspecies or species criterion was met. For review of species concepts, see Reeves et al. (2004), Orr and Coyne (2004), de Queiroz (2007), Perrin (2009) and Taylor et al. (2017 a,b). Nomenclatural papers that propose new names are to be registered with ZooBank and publications should include required information including designation of holotype or syntype specimen(s), identification of the repository of the specimen(s), and explicit identification of the taxon as new by inclusion of a term such as "ssp. nov.", "new species", etc. See the Committee's [Taxonomic Guide for Authors](#) for more information. The Committee also recommends authors acquaint themselves with the International Code of Zoological Nomenclature (ICZN 1999) and supply the information required under the Code when proposing new species or subspecies. Some examples of recent species or subspecies descriptions and redescrptions providing the necessary information can be seen at: Yamada et al. (2019), Rosel et al. (2021), Costa et al. (2023).

The Committee omits some described species and subspecies because of concern about their biological distinctness; reservations are given in the [narrative](#) below. In addition, published proposals for taxonomically distinct units that do not provide names and instead refer simply to

**Commented [Anon1]:** This is to be hyperlinked to the new document "Taxonomic guide for Authors."

'un-named subspecies' or 'un-named species' are not included in the official list but are provided [here](#). Authors are encouraged to provide formal descriptions following ICZN rules based on robust data showing an appropriate subspecies or species criterion is met.

Common names are arbitrary and change with time and place; one or two frequently used names in English and/or a range language are given here. Additional English common names and common names in French, Spanish, Russian and other languages are available at [www.marinespecies.org/cetacea/](http://www.marinespecies.org/cetacea/). The Committee does not arbitrate common names. In general, we aim to maintain established names currently in use whenever possible, use names suggested by the authors of new species or subspecies descriptions if present, and recognize multiple common names when multiple names are commonly used.

Species are listed in alphabetical order within families.

Corrections and comments should be directed to the Committee on Taxonomy ([patricia.rosel@noaa.gov](mailto:patricia.rosel@noaa.gov)). Divergent opinions by members of the Committee on particular taxonomic questions are given in the footnotes.

## **List of Marine Mammal Species and Subspecies**

### **Order Carnivora**

#### **Family Ursidae**

*Ursus maritimus* Linnaeus 1758. Polar bear

#### **Family Mustelidae**

*Enhydra lutris* (Linnaeus, 1758). Sea otter

- *E. l. kenyoni* Wilson, 1991. Eastern sea otter
- *E. l. lutris* (Linnaeus, 1758). Western sea otter
- *E. l. nereis* (Merriam, 1904). Southern sea otter

*Lontra felina* (Molina, 1782). Chungungo, marine otter

*Neovison macrodon* (Prentis, 1903). Sea mink (extinct)

### **PINNIPEDIA (eared seals, sea lions, walrus, earless seals; 35 species, of which 2 extinct)**

#### **Family Otariidae (eared seals and sea lions; 15 species, of which 1 extinct)**

*Arctocephalus australis* (Zimmermann, 1783). South American fur seal

*Arctocephalus forsteri* (Lesson, 1828). Long-nosed fur seal, New Zealand fur seal

*Arctocephalus galapagoensis* Heller, 1904. Galapagos fur seal

*Arctocephalus gazella* (Peters, 1875). Antarctic fur seal

*Arctocephalus philippii* (Peters, 1866). Juan Fernandez fur seal

- *A. p. philippii* (Peters, 1866). Juan Fernandez fur seal
- *A. p. townsendi* (Merriam, 1897). Guadalupe fur seal

*Arctocephalus pusillus* (Schreber, 1775). Cape fur seal

- *A. p. pusillus* (Schreber, 1775). Cape fur seal
- *A. p. doriferus* Wood Jones, 1925. Australian fur seal

*Arctocephalus tropicalis* (Gray, 1872). Subantarctic fur seal

*Callorhinus ursinus* (Linnaeus, 1758). Northern fur seal

*Eumetopias jubatus* (Schreber, 1776). Steller sea lion, northern sea lion

- *E. j. jubatus* (Schreber, 1776). Western Steller sea lion
- *E. j. monteriensis* (Gray, 1859). Loughlin's Steller sea lion

*Neophoca cinerea* (Peron, 1816). Australian sea lion

*Otaria byronia* (Blainville, 1820). South American sea lion

*Phocarctos hookeri* (Gray, 1844). New Zealand sea lion, Hooker's sea lion

*Zalophus californianus* (Lesson, 1828). California sea lion

*Zalophus japonicus* (Peters, 1866). Japanese sea lion (extinct)

*Zalophus wollebaeki* Sivertsen, 1953. Galapagos sea lion

#### **Family Odobenidae**

*Odobenus rosmarus* (Linnaeus, 1758). Walrus

- *O. r. divergens* (Illiger, 1815). Pacific walrus
- *O. r. rosmarus* (Linnaeus, 1758). Atlantic walrus

#### **Family Phocidae (earless seals; 19 species, of which 1 extinct)**

*Cystophora cristata* (Erxleben, 1777). Hooded seal

*Erignathus barbatus* (Erxleben, 1777). Bearded seal

- *E. b. barbatus* (Erxleben, 1777). Atlantic bearded seal
- *E. b. nauticus* (Pallas, 1881). Pacific bearded seal

*Halichoerus grypus* (Fabricius, 1791). Gray seal

- *H. g. grypus* (Fabricius, 1791). Baltic gray seal
- *H. g. atlantica*. Nehring, 1866. Atlantic gray seal

*Histiophoca fasciata* (Zimmerman, 1783). Ribbon seal

*Hydrurga leptonyx* (Blainville, 1820). Leopard seal

*Leptonychotes weddellii* (Lesson, 1826). Weddell seal

*Lobodon carcinophaga* (Hombron and Jacquinot, 1842). Crabeater seal

*Mirounga leonina* (Linnaeus, 1758). Southern elephant seal

*Mirounga angustirostris* (Gill, 1866). Northern elephant seal

*Monachus monachus* (Hermann, 1779). Mediterranean monk seal

*Neomonachus tropicalis* (Gray, 1850). Caribbean monk seal, West Indian monk seal (extinct)

*Neomonachus schauinslandi* (Matschie, 1905). Hawaiian monk seal

*Ommatophoca rossii* Gray, 1844. Ross seal

*Pagophilus groenlandicus* (Erxleben, 1777). Harp seal

*Phoca vitulina* Linnaeus, 1758. Harbor seal, common seal

- *P. v. vitulina* Linnaeus, 1758. Atlantic harbor seal
- *P. v. mellonae* Douutt, 1942. Ungava harbor seal
- *P. v. richardii* (Gray, 1864). Pacific harbor seal

*Phoca largha* Pallas, 1811. Spotted seal, largha seal

*Pusa hispida* (Schreber, 1775). Ringed seal

- *P. h. hispida* (Schreber, 1775). Arctic Ringed seal
- *P. h. botnica* (Gmelin, 1788). Baltic ringed seal
- *P. h. ochotensis* (Pallas, 1811). Okhotsk ringed seal
- *P. h. ladogensis* (Nordquist, 1889). Lake Ladoga seal

- *P. h. saimensis* (Nordquist, 1889). Saima seal

*Pusa caspica* (Gmelin, 1788). Caspian seal

*Pusa sibirica* (Gmelin, 1788). Baikal seal

## **Order ARTIODACTYLA<sup>1</sup> (artiodactyls and cetaceans)**

### **Infraorder CETACEA (cetaceans; 94 species, of which 1 possibly extinct)**

#### **MYSTICETI (baleen whales, 15 species)**

##### **Family Balaenidae (right whales, 4 species)<sup>2</sup>**

*Balaena mysticetus* Linnaeus, 1758. Bowhead whale, Greenland whale

*Eubalaena glacialis* (Müller, 1776). North Atlantic right whale

*Eubalaena japonica* (Lacépède, 1818). North Pacific right whale

*Eubalaena australis* (Desmoulins, 1822). Southern right whale

##### **Family Neobalaenidae**

*Caperea marginata* (Gray, 1846). Pygmy right whale

##### **Family Eschrichtiidae**

*Eschrichtius robustus* (Lilljeborg, 1861). Gray whale

##### **Family Balaenopteridae (rorquals, 9 species)**

*Balaenoptera acutorostrata* Lacépède, 1804. Common minke whale

- *B. a. acutorostrata* Lacépède, 1804. North Atlantic minke whale
- *B. a. scammoni* Deméré, 1986. North Pacific minke whale

*Balaenoptera bonaerensis* Burmeister, 1867. Antarctic minke whale

*Balaenoptera borealis* Lesson, 1828. Sei whale

- *B. b. borealis* Lesson, 1828. Northern sei whale
- *B. b. schlegelii* (Flower, 1865). Southern sei whale

*Balaenoptera edeni* Anderson, 1879. Bryde's whale

- *B. e. brydei* Olsen, 1913. Bryde's whale
- *B. e. edeni* Anderson, 1879. Eden's whale

*Balaenoptera musculus* (Linnaeus, 1758). Blue whale

- *B. m. musculus* (Linnaeus, 1758). Northern blue whale
- *B. m. intermedia* Burmeister, 1871. Antarctic blue whale
- *B. m. indica* Blyth, 1859. Northern Indian Ocean blue whale
- *B. m. brevicauda* Ichihara, 1966. Pygmy blue whale

*Balaenoptera omurai* Wada, Oishi and Yamada, 2003. Omura's whale

*Balaenoptera physalus* (Linnaeus, 1758). Fin whale

- *B. p. physalus* (Linnaeus, 1758). North Atlantic fin whale
- *B. p. quoyi* (Fischer, 1829). Southern fin whale
- *B. p. velifera* Cope in Scammon, 1869. North Pacific fin whale

*Balaenoptera ricei* Rosel, Wilcox, Yamada and Mullin, 2021. Rice's whale

*Megaptera novaeangliae* (Borowski, 1781). Humpback whale

- *M. n. australis* (Lesson, 1828). Southern humpback whale
- *M. n. kuzira* (Gray, 1850). North Pacific humpback whale
- *M. n. novaeangliae* (Borowski, 1781). North Atlantic humpback whale

**ODONTOCETI (toothed whales, dolphins and porpoises: 79 species, of which one possibly extinct)**

#### **Family Physeteridae**

*Physeter macrocephalus* Linnaeus, 1758. Sperm whale, cachalot

#### **Family Kogiidae**

*Kogia breviceps* (Blainville, 1838). Pygmy sperm whale

*Kogia sima* (Owen, 1866). Dwarf sperm whale

#### **Family Ziphiidae (beaked whales, 24 species)**

*Berardius arnuxii* Duvernoy, 1851. Arnoux's beaked whale

*Berardius bairdii* Stejneger, 1883. Baird's beaked whale

*Berardius minimus* Yamada, Kitamura and Matsuishi, 2019. Sato's beaked whale

*Hyperoodon ampullatus* (Forster in Kalm, 1770). Northern bottlenose whale  
*Hyperoodon planifrons* Flower, 1882. Southern bottlenose whale  
*Indopacetus pacificus* (Longman, 1926). Longman's beaked whale, tropical bottlenose whale  
*Mesoplodon bidens* (Sowerby, 1804). Sowerby's beaked whale  
*Mesoplodon bowdoini* Andrews, 1908. Andrews' beaked whale  
*Mesoplodon carlhubbsi* Moore, 1963. Hubbs' beaked whale  
*Mesoplodon eueu* Carroll et al. 2021<sup>4</sup>. Ramari's beaked whale  
*Mesoplodon densirostris* (Desmarest, 1817). Blainville's beaked whale  
*Mesoplodon europaeus* (Gervais, 1855). Gervais' beaked whale  
*Mesoplodon ginkgodens* Nishiwaki and Kamiya, 1958. Ginkgo-toothed beaked whale  
*Mesoplodon grayi* von Haast, 1876. Gray's beaked whale  
*Mesoplodon hectori* (Gray, 1871). Hector's beaked whale  
*Mesoplodon hotaula* Deraniyagala, 1963. Deraniyagala's beaked whale  
*Mesoplodon layardii* (Gray, 1865). Strap-toothed beaked whale, Layard's beaked whale  
*Mesoplodon mirus* True, 1913. True's beaked whale  
*Mesoplodon perrini* Dalebout, Mead, Baker, Baker and van Helden, 2002. Perrin's beaked whale  
*Mesoplodon peruvianus* Reyes, Mead and Van Waerebeek, 1991. Pygmy beaked whale  
*Mesoplodon stejnegeri* True, 1885. Stejneger's beaked whale  
*Mesoplodon traversii* (Gray, 1874). Spade-toothed whale  
*Tasmacetus shepherdi* Oliver, 1937. Shepherd's beaked whale, Tasman beaked whale  
*Ziphius cavirostris* G. Cuvier, 1823. Cuvier's beaked whale, goose-beaked whale

#### **Family Platanistidae**

*Platanista gangetica* (Lebeck, 1801). Ganges river dolphin, susu

*Platanista minor* Owen, 1853. Indus river dolphin, bhulan

#### **Family Iniidae**

*Inia geoffrensis* (Blainville, 1817). Amazon river dolphin

- *I. g. boliviensis* (d'Orbigny, 1834). Bolivian bufeo
- *I. g. geoffrensis* (Blainville, 1817). Common boto

#### **Family Lipotidae**

*Lipotes vexillifer* Miller, 1918. Baiji, Yangtze river dolphin (possibly extinct)

#### **Family Pontoporiidae**

*Pontoporia blainvillei* (Gervais and d'Orbigny, 1844). Franciscana, toninha

#### **Family Monodontidae**

*Delphinapterus leucas* (Pallas, 1776). Beluga, white whale

*Monodon monoceros* Linnaeus, 1758. Narwhal

#### **Family Delphinidae (38 species)**

*Cephalorhynchus commersonii* (Lacépède, 1804). Commerson's dolphin

- *C. c. commersonii* (Lacépède, 1804). Commerson's dolphin
- *C. c. kerguelenensis* Robineau, Goodall, Pichler and C. S. Baker, 2007. Kerguelen dolphin

*Cephalorhynchus eutropia* (Gray, 1846). Chilean dolphin

*Cephalorhynchus heavisidii* (Gray, 1828). Heaviside's dolphin

*Cephalorhynchus hectori* (Van Beneden, 1881). Hector's dolphin

- *C. h. hectori* (Van Beneden, 1881). South Island Hector's dolphin
- *C. h. maui* A. Baker, Smith and Pichler, 2002. Māui dolphin, North Island Hector's dolphin

*Delphinus delphis* Linnaeus, 1758. Common dolphin

- *D. d. delphis* Linnaeus, 1758. Common dolphin
- *D. d. bairdii* Dall, 1873. Eastern North Pacific long-beaked common dolphin<sup>3</sup>
- *D. d. ponticus* Barabash, 1935. Black Sea common dolphin



- *D. d. tropicalis* van Bree, 1971. Indo-Pacific common dolphin

*Feresa attenuata* Gray, 1874. Pygmy killer whale

*Globicephala macrorhynchus* Gray, 1846. Short-finned pilot whale

*Globicephala melas* (Traill, 1809). Long-finned pilot whale

- *G. m. edwardii* (A. Smith, 1834). Southern long-finned pilot whale
- *G. m. melas* (Traill, 1809). North Atlantic long-finned pilot whale

*Grampus griseus* (G. Cuvier, 1812). Risso's dolphin, grampus

*Lagenodelphis hosei* Fraser, 1956. Fraser's dolphin

*Lagenorhynchus acutus* (Gray, 1828). Atlantic white-sided dolphin

*Lagenorhynchus albirostris* (Gray, 1846). White-beaked dolphin

*Lagenorhynchus australis* (Peale, 1849). Peale's dolphin

*Lagenorhynchus cruciger* (Quoy and Gaimard, 1824). Hourglass dolphin

*Lagenorhynchus obliquidens* Gill, 1865. Pacific white-sided dolphin

*Lagenorhynchus obscurus* (Gray, 1828). Dusky dolphin

- *L. o. fitzroyi* (Waterhouse, 1838). Fitzroy's dolphin
- *L. o. obscurus* (Gray, 1828). African dusky dolphin
- *L. o. posidonia* (Philippi, 1893). Peruvian/Chilean dusky dolphin

*Lissodelphis borealis* (Peale, 1849). Northern right-whale dolphin

*Lissodelphis peronii* (Lacépède, 1804). Southern right-whale dolphin

*Orcaella brevirostris* (Owen in Gray, 1866). Irrawaddy dolphin, pesut

*Orcaella heinsohni* Beasley, Robertson and Arnold, 2005. Australian snubfin dolphin

*Orcinus orca* (Linnaeus, 1758). Killer whale, orca

*Peponocephala electra* (Gray, 1846). Melon-headed whale

*Pseudorca crassidens* (Owen, 1846). False killer whale

*Sousa teuszii* (Kükenthal, 1892). Atlantic humpback dolphin

*Sousa chinensis* (Osbeck, 1765). Indo-Pacific humpback dolphin

- *S. c. chinensis* (Osbeck, 1765). Chinese humpback dolphin
- *S. c. taiwanensis* Wang, Yang and Hung, 2015. Taiwanese humpback dolphin

*Sousa plumbea* (G. Cuvier, 1829). Indian Ocean humpback dolphin

*Sousa sahalensis* Jefferson and Rosenbaum, 2014. Australian humpback dolphin

*Sotalia fluviatilis* (Gervais and Deville in Gervais, 1853). Tucuxi

*Sotalia guianensis* (P.J. Van Beneden, 1864). Guiana dolphin, costero

*Stenella attenuata* (Gray, 1846). Pantropical spotted dolphin

- *S. a. attenuata* (Gray, 1846). Offshore pantropical spotted dolphin
- *S. a. graffmani* (Lönnerberg, 1934). Coastal pantropical spotted dolphin

*Stenella clymene* (Gray, 1850). Clymene dolphin

*Stenella coeruleoalba* (Meyen, 1833). Striped dolphin

*Stenella frontalis* (G. Cuvier, 1829). Atlantic spotted dolphin

*Stenella longirostris* (Gray, 1828). Spinner dolphin

- *S. l. centroamericana* Perrin, 1990. Central American spinner dolphin
- *S. l. longirostris* (Gray, 1828). Gray's spinner dolphin
- *S. l. orientalis* Perrin, 1990. Eastern spinner dolphin
- *S. l. roseiventris* (Wagner, 1846). Dwarf spinner dolphin

*Steno bredanensis* (Lesson, 1828). Rough-toothed dolphin

*Tursiops aduncus* (Ehrenberg, 1832). Indo-Pacific bottlenose dolphin

*Tursiops erebennus* (Cope, 1865). Tamanend's bottlenose dolphin

*Tursiops truncatus* (Montagu, 1821). Common bottlenose dolphin

- *T. t. gephyreus* Lahille, 1908. Lahille's bottlenose dolphin
- *T. t. nuuanu* Andrews, 1911. Eastern Tropical Pacific bottlenose dolphin
- *T. t. ponticus* Barabash-Nikiforov, 1940. Black Sea bottlenose dolphin
- *T. t. truncatus* (Montagu, 1821). Common bottlenose dolphin

### **Family Phocoenidae (porpoises, 7 species)**

*Neophocaena phocaenoides* (G. Cuvier, 1829). Indo-Pacific finless porpoise

*Neophocaena asiaeorientalis* (Pilleri and Gahr, 1972). Narrow-ridged finless porpoise

- *N. a. asiaeorientalis* (Pilleri and Gahr, 1972). Yangtze finless porpoise
- *N. a. sunameri* Pilleri and Gahr, 1975. East Asian finless porpoise

*Phocoena dioptrica* Lahille, 1912. Spectacled porpoise

*Phocoena phocoena* (Linnaeus, 1758). Harbor porpoise

- *P. p. phocoena* (Linnaeus, 1758). Atlantic harbor porpoise
- *P. p. vomerina* (Gill, 1865). Pacific harbor porpoise
- *P. p. relicta* Abel, 1905. Black Sea harbor porpoise

*Phocoena sinus* Norris and McFarland, 1958. Vaquita, Gulf of California harbor porpoise

*Phocoena spinipinnis* Burmeister, 1865. Burmeister's porpoise

*Phocoenoides dalli* (True, 1885). Dall's porpoise

- *P. d. dalli* (True, 1885). *dalli*-type Dall's porpoise
- *P. d. truei* Andrews, 1911. *truei*-type Dall's porpoise

### **ORDER SIRENIA (sirenians, 5 species, of which 1 extinct)**

#### **Family Trichechidae**

*Trichechus inunguis* (Natterer, 1883). Amazonian manatee

*Trichechus manatus* Linnaeus, 1758. West Indian manatee

- *T. m. latirostris* (Harlan, 1824). Florida manatee
- *T. m. manatus* Linnaeus, 1758. Antillean manatee

*Trichechus senegalensis* Link, 1795. West African manatee, African manatee

#### **Family Dugongidae**

*Dugong dugon* (Müller, 1776). Dugong

*Hydrodamalis gigas* (Zimmerman, 1780). Steller's sea cow (extinct)

#### **Footnotes (dissenting opinions):**

1 Use of Order Cetartiodactyla (artiodactyls and cetaceans) is favored by some evolutionary mammalogists working with molecular data. Some others, including many marine mammalogists and paleontologists, favor retention of Order Cetacea in the interest of taxonomic stability, despite rendering the retained Order Artiodactyla paraphyletic.

2 (from D. Rice) Baker et al. (2003) hold that there is no evidence that would support the classification of the right whales as more than a single biological species. [The three species are here recognized as phylogenetic species.]

3 (from T. Jefferson) The List should retain the eastern North Pacific (ENP) long-beaked common dolphin as a separate species of *Delphinus*, *D. bairdii* (as in Banks and Brownell 1969). The evidence for species status is published and is considerable, especially the molecular differences. Notwithstanding issues related to poor sampling from areas of potential sympatry further south, the balance of the evidence seems to support a lack of interbreeding throughout a very large region of overlap, along with clear ecological differences. In this view, the error by Heyning and Perrin (1994) was not in splitting out the ENP long-beaked common dolphins as a distinct species, but in assuming that the name *D. capensis* applied there and that all long-beaked populations of *Delphinus* belonged to *D. capensis*.

<sup>4</sup>The full authority is: Carroll, McGowen, McCarthy, Marx, Aguilar, Dalebout, Dreyer, Gaggiotti, Hansen, van Helden, Onoufriou, Baird, Baker, Berrow, Cholewiak, Claridge, Constantine, Davison, Eira, Fordyce, Gatesy, Greg Hofmeyr, Martin, Mead, Mignucci-Giannoni, Morin, Reyes, Rogan, Rosso, Silva, Springer, Steel and Tange Olsen, 2021

## **Narrative on Taxonomy**

Based on molecular and morphological data, the cetaceans fall firmly within the artiodactyl clade with hippopotamuses being the sister taxon to cetaceans (e.g., Geisler and Uhen, 2005, Spaulding et al. 2009) and Montgelard et al. (1997) proposed the taxon Cetartiodactyla, pooling even-toed terrestrial mammals and all cetaceans into a single Order. The Taxonomy Committee followed this convention, with some reservation, when it created the first list of marine mammal species and subspecies in 2009, and the term Cetartiodactyla has been widely used. Recently, Prothero et al. (2022) make a strong argument that the term Cetartiodactyla misrepresents taxonomic relationships as it implies cetaceans and artiodactyls are of equal rank, when, in fact, cetaceans are clearly nested within the Artiodactyla. We agree with Prothero et al. (2022) and revert to the use of Order Artiodactyla to include all cetacean taxa (living and extinct), and provisionally rank Cetacea as an Infraorder. While this change will likely cause some initial confusion, it more accurately reflects the placement of cetaceans in the taxonomic hierarchy. We continue to maintain Mysticeti and Odontoceti provisionally as unranked taxa (recognizing that the classification within Cetartiodactyla remains partially unresolved — e.g., see Spaulding et al. (2009), Price et al. (2005), Agnarsson and May-Collado (2008)).<sup>1</sup>

Below the rank of order, we list only families, genera, species and subspecies, omitting superfamilies, subfamilies and taxa of other ranks. Consistent with the use of Cetacea as an un-

ranked unit, we also include the un-ranked taxon Pinnipedia. Morphological (Wyss and Flynn, 1993; Berta and Wyss, 1994) and molecular (Higdon et al. 2007; Fulton and Strobeck, 2010, Nyakatura and Bininda-Emonds 2012) analyses provide strong support for pinniped monophyly and hence inclusion of Pinnipedia as an un-ranked taxonomic unit. However, there is continued debate on this subject as Koretsky et al. (2016) argued the case for a diphyletic origin for pinnipeds.

### Pinnipeds

For pinnipeds, we previously followed Berta and Churchill (2012). To avoid issues of paraphyly, these authors proposed that, based on molecular and morphological data, the genus *Arctocephalus* be limited to *Arctocephalus pusillus*, the type species of the genus *Arctocephalus*, and transferred the remaining 'Arctocephalus' species (i.e., *A. australis*, *A. galapagoensis*, *A. gazella*, *A. philippii* and *A. tropicalis*) to *Arctophoca* Peters, 1866. However, Nyakatura and Bininda-Emonds (2012) compiled a new supertree of the Carnivora and concluded that this usage of *Arctophoca* may be premature because of remaining uncertainty about phylogenetic relationships, and we return provisionally to use of *Arctocephalus* for all the southern fur seals.

Three subspecies of *A. australis* were proposed by Berta and Churchill (2012) and formerly listed here: *A. a. australis*, *A. a. forsteri* and *A. a. gracilis*. However, Oliveira and Brownell (2014) synonymized *A. a. gracilis* with *A. a. australis*. The super-tree analysis by Nyakatura and Bininda-Emonds (2012) accords with the phylogenetic analysis of Higdon et al. (2007), suggesting that the New Zealand fur seal should be recognized as a full species, *A. forsteri*. Two subspecies of *A. philippii* are thought to be biologically distinct: *A. p. philippii* and *A. p. townsendi*, although small sample sizes and a small number of genes sampled are concerns. Two subspecies of *Eumetopias* are supported largely on molecular genetic data, which is also the case for recognition of California, Japanese and Galapagos sea lions as separate species. Following a proposal by Gardner and Robbins (1999), the International Code of Zoological Nomenclature (ICZN) placed the name *O. byronia* on its Official List of Specific Names in Zoology. Brunner (2004) also advised use of *Otaria byronia* (Blainville, 1820) over *O. flavescens* (Shaw, 1800) and Webber (2014) provides a summary of why *O. flavescens* is a *nomen dubium* and should not be used under the provisions of the International Code of Zoological Nomenclature (ICZN 1999). However, many South American scientists continue to use *O. flavescens*.

Lindqvist et al. (2009) concluded that a purported third subspecies of the walrus *Odobenus rosmarus laptevi* is not warranted.

Recent molecular genetic analyses indicate that *Phoca vitulina concolor* is paraphyletic and this along with lack of morphological differentiation suggests that the western Atlantic subspecies is not supportable; *P. v. vitulina* is considered here to apply to all Atlantic harbor seals. Within the North Pacific, until the subspecies limits of various populations are assessed, only a single subspecies is recognized, *Phoca vitulina richardii*. Placement of the ringed seal, Caspian seal and Baikal seal has alternated between the genera *Phoca* and *Pusa*. We accept Rice's (1998) use of *Pusa* as the correct classification. Scheel et al. (2014) found the Caribbean and Hawaiian monk seals to be more closely related molecularly and morphologically to each other than either is to the Mediterranean monk seal and created the new genus *Neomonachus* for the two species.

The long-lost holotype skull of the gray seal *Halichoerus grypus* has been rediscovered and shown by DNA analysis to hail from the Baltic rather than from Greenland as previously thought (Olsen et al. 2016). Consequently, the nominate subspecies *H. g. grypus* is the Baltic gray seal, *H. g. macrorhynchus* falls into synonymy, and the authors have resurrected *H. g. atlantica* to apply to the Atlantic subspecies.

#### Polar Bears and Otters

Derocher and Stirling (1998) argued convincingly that patterns of variation in the polar bear do not support recognition of subspecies.

The use of *Lontra* rather than *Lutra* for the marine otter follows Larivière (1998) in recognizing the otters of North and South America as a monophyletic taxon distinct from the otters of Eurasia.

#### Baleen Whales

In the mysticete cetaceans, molecular evidence strongly supports the recognition of three separate phylogenetic species of right whales (Rosenbaum et al. 2000; Gaines et al. 2005). In addition, the genus *Eubalaena* (rather than *Balaena* as in Rice, 1998) is retained for the right whales as recommended by the Scientific Committee of the International Whaling Commission (IWC, 2001)<sup>2</sup>.

New fossil evidence suggests that *Caperea marginata* may be a member of the family Cetotheriidae (Fordyce and Marx 2012; Marx and Fordyce 2015; Marx and Fordyce 2016). Neobalaenidae is retained here provisionally.

Jackson et al. (2014) have recognized three subspecies of the humpback whale based on mitochondrial and nuclear DNA relationships and distribution: *Megaptera novaeangliae kuzira* (North Pacific), *M. n. novaeangliae* (North Atlantic) and *M. n. australis* (Southern Hemisphere).

There have been a variety of studies (McDonald et al. 2006, Branch et al. 2007, LeDuc et al. 2007, 2017, Torres-Florez et al. 2014, Buchan et al. 2018, Leslie et al. 2021) that have provided evidence suggesting blue whales in the eastern South Pacific differ from blue whales elsewhere and may represent a separate subspecies but none have provided a formal subspecies description supplying the information required under the International Code of Zoological Nomenclature for proposing new species or subspecies. Khalif (2020) summarizes information available on the population in the eastern South Pacific, designates an identifiable specimen and type location, and provides a subspecies name. However, the subspecies description is incomplete and, importantly, is diagnosed by bibliographic references alone. This is minimally sufficient to make the name available (see Code, Art. 13.1.2), but no attempt seems to have been made to collect morphological or genetic evidence, despite the ready availability of a skull and nearly complete skeleton. This is poor practice and contrary to Recommendation 13A. We therefore regard the name *B. musculus chilensis* a *nomen dubium* and do not recognize it until these faults are remedied. Khalif (2021) also describes a new blue whale subspecies *B. musculus arabica* from the Arabian Sea. This work fails to acknowledge that more than one type of blue whale exists in

northern Indian Ocean making it impossible to link the new blue whale song described by Cerchio et al. (2020) from this region to the blue whale skeleton from Kuwait. The subspecies is diagnosed only by bibliographic references to characterizations of its song and acoustics alone does not constitute sufficient evidence for taxonomic separation. We therefore regard the name *B. musculus arabica* a *nomen dubium* and do not recognize it.

The Bryde's-like whales in the Gulf of Mexico have now been described as a new species of medium-sized balaenopterid, *Balaenoptera ricei*, by Rosel et al. (2021), with the common name Rice's whale. The remaining Bryde's whales are provisionally considered to comprise a single species, *Balaenoptera edeni*, following the usage of Kato and Perrin (2009) and Kershaw et al. (2013). However, Luksenburg et al. (2015) and Wada et al. (2003) accepted two species. Some workers recognize *B. edeni* as including only the small-form coastal Bryde's whales of the western Pacific and Indian Oceans, using *B. brydei* for the globally distributed and larger, more oceanic form (Sasaki et al. 2006). Kato and Perrin (2009) and Kershaw et al. (2013) considered these more likely to be distinct at the subspecific level (although arguably at the species level), and they are included here provisionally as such. *Balaenoptera omurai* was described by Wada et al. (2003). It was previously confounded with the Bryde's whale and has been confirmed as having a separate and ancient lineage (Sasaki et al. 2006).

To further resolve the taxonomy and systematics of all the medium-sized baleen whales (*B. ricei*, *B. edeni edeni*, *B. e. brydei*, and including *B. borealis*) it is critical that a neotype be designated for *B. brydei* and that genetic data be obtained from that designated neotype. In addition, genetic data should also be obtained from the holotype of *B. edeni* to validate the identity of museum specimens as well as samples collected from stranded and biopsied individuals. Finally, a comprehensive phylogenetic analysis using nuclear markers of all taxa in this complex with sufficient samples and range-wide geographic coverage is also needed.

Clarke (2004) proposed recognition of a pygmy form of the fin whale as a subspecies, based on distribution, size and coloration. He resurrected the synonym *patachonica* Burmeister, 1865 to apply to the subspecies: *B. physalus patachonica*. However, Pérez-Alvarez et al. (2021) conducted a phylogenetic analysis of fin whales and found no support for differentiation among Southern Hemisphere fin whales, recommending that the name *patachonica* Burmeister, 1865 be synonymized with *B. physalus quoyi* and the latter refer to all Southern Hemisphere fin whales.

Archer et al. (2019) provided genetic evidence at both mitochondrial and nuclear genetic markers that fin whales in the North Pacific are significantly differentiated from those in the North Atlantic and the Southern Hemisphere. They proposed recognition of the North Pacific fin whale as a separate subspecies, *B. physalus velifera*.

#### Toothed Whales

In the odontocetes, *Mesoplodon traversii* (spade-toothed whale) was recognized as the senior synonym for *M. bahamondi* (Bahamonde's beaked whale) (van Helden et al. 2002). The first complete specimen was recently described from a stranding on the North Island of New Zealand (Thompson et al. 2012). *Mesoplodon perrini* was described by Dalebout et al. (2002). Dalebout et al. (2014) resurrected *Mesoplodon hotaula* Deraniyagala, 1963, a species closely similar to *M.*

*ginkgodens*. Using morphological and genomic evidence, Carrol et al. (2021) split *M. mirus*, identifying those in the Southern Hemisphere as *M. eueu*.

Yamada et al. (2019) provided morphological evidence, supported by genetic data (Morin et al. 2018, Kitamura et al. 2013), recognizing a new species of *Berardius* in the North Pacific, *Berardius minimus*.

The South Asian river dolphins (genus *Platanista*) have a long history of taxonomic revision, oscillating between being recognized as two species, *Platanista gangetica* (Lebeck, 1801) and *P. minor* Owen, 1853, and two subspecies of *Platanista gangetica*. The Taxonomy Committee, when it first created the taxonomy list, followed Rice's (1998) treatment of *Platanista* and recognized a single species with two subspecies, *P. g. gangetica* and *P. g. minor*. Recently, Braulik et al. (2014, 2021) have performed a morphological analysis of the two taxa, coupled with an analysis of mitochondrial DNA and recommended species-level status for the two subspecies. Individually, the support provided from these two lines of evidence was relatively weak: a single, but significant, diagnostic difference in the frontal bones of the skull, and five fixed differences and no shared mtDNA haplotypes, but a level of genetic divergence that does not exceed the species threshold identified by Taylor et al. (2017a). The combined evidence for diagnosability in the morphological and genetic lines of evidence, the significant difference in growth curves between the two taxa, and their estimated divergence time of ~0.55 million years, led the majority of committee members to accept the proposed elevation to species. However, the unavailability of samples from the upper reaches of the rivers where the two taxa may have last been in contact impedes a complete understanding of the evolutionary history of the two taxa and this posed some concern for a few members when evaluating the proposal to elevate the subspecies to species.

We recognize one species of *Inia* with two subspecies: *I. geoffrensis geoffrensis* (Blainville, 1817) and *I. g. boliviensis* (d'Orbigny, 1834). *I. g. boliviensis* is found in the Bolivian Amazon basin and overlaps with other *Inia* in all morphological characters (da Silva, 1994; Ruiz-García et al. 2006). However, molecular genetic evidence from mitochondrial DNA (mtDNA) and nuclear introns (Banguera-Hinestroza et al. 2002; Ruiz-García et al. 2008) suggested these two subspecies are on separate evolutionary trajectories and deserve recognition as phylogenetic species. Hollatz et al. (2011) estimated that *I. g. boliviensis* has been reproductively isolated for 3.2 million years. However, these studies used geographically disparate samples, with a significant gap in sampling of *Inia* habitat between the Teotonio Rapids, hypothesized to be the barrier to gene flow for *I. g. boliviensis*, and the samples used to represent *I. g. geoffrensis*. Gravena et al. (2014), with much better sampling of the Madeira River system, including samples from immediately above and below the Teotonio Rapids, found that in fact these rapids did not appear to obstruct gene flow on an evolutionary scale. Thus, the conclusion that the Bolivian *Inia* above the Teotonio Rapids possessed unique mtDNA (a major line of evidence for recognizing species-level distinctness of *I. boliviensis*) was not supported by more extensive sampling (i.e., the initial appearance of separation was an artifact of poor sampling). Therefore, only the subspecies *Inia geoffrensis boliviensis* is retained, although given the newer evidence for a lack of isolation of the dolphins above the Teotonio Rapids (Gravena et al. 2014) and the lack of robust sampling, the strength for subspecies status is weakened. Another new species, *Inia araguaiaensis*, was described by Hrbek et al. (2014) from the Araguaia River, which is not



connected to the Amazon and Siciliano et al. (2016) extended the known range of *I. araguaiaensis* further north to Marajó Bay, Brazil through mtDNA analysis of three stranded specimens. Hrbek et al. (2014) only examined samples from two extremes of the distribution of *Inia*, so is it unclear if the molecular differences observed represented real species-level separation or were due to sampling from two locations separated by a large distance. Diagnostic osteological differences were also reported (Hrbek et al. 2014). However, because this study was based on the examination of very few specimens (only 2 for the new species and only 9 for *I. geoffrensis*) and did not account for effects of sexual dimorphism within the species, the authors' conclusions are not convincing. In light of these arguments, the species here remains unlisted provisionally. Hrbek et al. (2014) did not recognize *I. g. humboldtiana*. Melo et al. (2021) examined acoustic evidence and found some suggestion of vocal differences, but vocalizations are known to be plastic in odontocetes. In addition, there was significant overlap in many of the measured parameters and classification based on the acoustic data exhibited an error rate near 30%, which is quite high for taxonomic decisions. Most recently, Cañizales (2020) and Emin-Lima et al. (2022) provide valuable new morphological data to evaluate aspects of the taxonomy within the genus *Inia*. Cañizales (2020) examined 20 skulls from Venezuela identified as representing *I. g. humboldtiana* and provided some evidence for differences between these samples and those of *I. g. geoffrensis*, but differences were slight, with overlap in several analyses. Emin-Lima et al. (2022) also focused on skull morphology and examined 46 skulls from distant locales within the Amazon basin and identified three morphological groups that largely corresponded with *I. g. geoffrensis*, *I. g. humboldtiana*, and *I. g. boliviensis*. They found no differences between *I. g. geoffrensis* and *I. araguaiaensis*.

Resolution of the taxonomic status of all *Inia* forms requires an integrated approach using multiple lines of evidence, improved sampling throughout the distribution, and a thorough review of all the evidence (morphological, genetic, ecological, etc.). In addition, proposed changes to taxonomy will require a detailed taxonomic treatment, must provide diagnoses, and review relevant nomenclature. Some examples of recent species or subspecies descriptions and redescrptions providing the necessary taxonomic treatment can be seen at: Yamada et al. (2019), Braulik et al. (2021), Rosel et al. (2021), Costa et al. (2023).

Previous editions of this list recognized two globally distributed species of common dolphins: the short-beaked common dolphin *D. delphis* and the long-beaked common dolphin *D. capensis* as proposed by Heyning and Perrin (1994). However, evidence that *D. capensis* thus considered is a polyphyletic taxon has been accumulating for some time. The long-beaked condition is apparently a convergent character state induced by regional ecology. In some regions, long-beaked common dolphins are genetically more closely related to short-beaked common dolphins than to long-beaked common dolphins in other regions (e.g., see Natoli et al. 2006). Cunha et al. (2015) summarized the relevant data and analyses, along with additional molecular data and analysis, and recommended that *Delphinus capensis* not be further used as proposed by Heyning and Perrin (1994). That recommendation is followed here. Cunha et al. (2015) noted that because the sympatric/parapatric long-beaked and short-beaked common dolphins off California in the eastern North Pacific (ENP) appear not to interbreed, perhaps the ENP long-beaked common dolphins might be recognized as a separate species *D. bairdii* Dall, 1873 (as advocated by Banks and Brownell (1969) based on the long-beaked condition). However, the molecular analysis for the ENP (Rosel et al. 1994) did not include common dolphins from the contiguous regions to the

south in the eastern tropical and eastern South Pacific. Pending a more complete global review and revision of the common dolphins, the long-beaked ENP form is considered here provisionally as a subspecies *D. delphis bairdii* following the usage of Hershkovitz (1966)<sup>3</sup>. *Delphinus capensis* Gray, 1828 from South Africa remains in the synonymy of *Delphinus delphis* Linnaeus, 1758. The long-beaked common dolphin of the Indian Ocean is retained as a subspecies *D. delphis tropicalis*.

Based on a combined analysis of molecular genetic and morphological data, Mendez et al. (2013) proposed recognizing four species of *Sousa* (the humpback dolphins): the previously here-listed *S. teuszii* and *S. chinensis* plus *S. plumbea* and a new un-named species from the waters of northern Australia that was originally proposed by Frère et al. (2008, 2011) based on molecular data. A drawback of the phylogenetic analyses by Mendez et al. (2013) is that there was only one sample from the area of supposed sympatry of *S. plumbea* and *S. chinensis* and very low coverage of the Indo-Malay region (n=5). The two species are listed here provisionally, pending the outcome of further analysis including more samples from those areas. The new species from Australia has been given the name *Sousa sahulensis* (Jefferson and Rosenbaum 2014).

The new subspecies *Sousa chinensis taiwanensis* was described by Wang et al. (2015).

The Burrunan dolphin *Tursiops australis*, described by Charlton-Robb et al. (2011), is not included here; its basis is questionable because of several potential problems: 1) the specimens were compared morphologically only with bottlenose dolphins from Australia; 2) despite the small sample sizes, the series overlapped in all metric characters and separation was possible only with multivariate analysis (which commonly resolves geographical forms within a species, e.g., see Perrin et al. (1999) and Perrin et al. (2011) for *Stenella longirostris* and *Tursiops truncatus*, respectively); 3) comparisons of external morphology and non-metric characters were made only with *T. truncatus*, to the exclusion of *T. aduncus*; and 4) support for important nodes in molecular trees suggesting phylogenetic separation was low. A rigorous re-evaluation of the relevant data and arguments is needed. Recently, Jedensjö et al. (2020) conducted a broader morphological comparison of *Tursiops* skulls from around Australia, including skulls of both *T. truncatus* and *T. aduncus* and their respective holotypes, and did not find support for the Burrunan bottlenose dolphin, *T. australis*, proposed by Charlton-Robb et al. (2011). Skulls previously classified as *T. australis* all fell well within the *T. truncatus* group. In contrast, Moura et al. (2020) placed *T. australis* within a *T. aduncus* clade based on a nuclear genetic phylogeny.

Based on morphological and genetic analyses, Costa et al. (2022) described a new species of *Tursiops* from the western North Atlantic, separating the long-recognized coastal and offshore morphotypes from the western North Atlantic into two species. The offshore morphotype remains *Tursiops truncatus*, while the coastal morphotype found in nearshore coastal waters as well as the bays and estuaries along the U.S. east coast was split out as the new species *T. erebennus* (Cope, 1865). Additional work is needed to identify the full geographic distribution of *T. erebennus* in the Gulf of Mexico and Caribbean.

Costa et al. (2023) evaluated morphological variation among three distinct morphotypes of *T. truncatus* in the eastern Pacific, the southern California/Mexico coastal form, the northern

temperate offshore form, and an eastern tropical Pacific (ETP) offshore form. Cranial morphology and body size separate the eastern tropical Pacific (ETP) offshore form from other Pacific *Tursiops*. The authors resurrected the name *T. t. nuuanu* Andrews, 1911 for this subspecies.

Costa et al. (2016), Wickert et al. (2016), and Hohl et al. (2020) have examined morphological characteristics of *Tursiops* specimens from the western South Atlantic. Costa et al. (2016) recommended subspecies status for the larger coastal form found between southern Brazil and northern Argentina, while both Wickert et al. (2016) and Hohl et al. (2020) have recommended species status. Costa et al. (2021) paired morphological and genetic data from samples collected from both forms in southern Brazil and found strong congruence between morphological and mitochondrial and nuclear divergence; genetic differentiation was also observed in Oliveira et al. (2019). In both genetic studies, however, evidence for shared mitochondrial haplotypes and possibly shared nuclear ancestry was found. We provisionally recognize the subspecies *T. t. gephyreus* Lahille 1908 (Lahille's bottlenose dolphin) for this larger coastal form. Improved geographic sampling and improved objectivity in hypothesis testing is needed to fully resolve the taxonomy of *Tursiops* in this region. In particular, further work is needed to better identify the distribution of the *gephyreus* form and determine its phylogenetic relationship to coastal and offshore *Tursiops* in the western North Atlantic and throughout Caribbean waters. Wang et al. (2021) showed *gephyreus* to be the correct trivial name for this taxon (as long as it continues to be recognized as a subspecies or species).

Recognition of the Black Sea bottlenose dolphin is now well-supported by molecular genetic data (Viaud-Martinez et al. 2008), as is the Black Sea common dolphin (Natoli et al. 2008).

Kinze (2018) proposed that the scientific name for *Tursiops truncatus* should be *Tursiops tursio* as *tursio* has priority over *truncatus*. The committee noted that resurrecting the name *T. tursio* for the bottlenose dolphin would upset long-standing usage and would cause considerable confusion in national and international treaties, agreements, etc. Therefore, the name *T. truncatus* should continue to be used for the common bottlenose dolphin to maintain stability.

The genus *Lagenorhynchus* is widely considered a polyphyletic taxon containing morphologically convergent species (Cipriano 1997, LeDuc et al. 1999, McGowen 2011, Banguera-Hinestroza et al. 2014, Vollmer et al. 2019). Vollmer et al. (2019) provided a detailed review of the relationships among members of the genus *Lagenorhynchus* and their relationship to *Cephalorhynchus* and *Lissodelphis*. Results continue to support the polyphyletic nature of the genus *Lagenorhynchus* and multiple morphological and genetic analyses indicate that *L. albirostris* and *L. acutus* are evolutionarily distinct from the other members of all three genera. Vollmer et al. (2019) recommended revision of the genus *sensu* LeDuc et al. (1999), placing *L. obscurus*, *L. obliquidens*, *L. australis* and *L. cruciger* into the genus *Sagmatias*, *L. acutus* into the genus *Leucopleurus* and leaving *L. albirostris* in a monotypic genus *Lagenorhynchus*. While a majority of committee members supported the changes proposed by Vollmer et al. (2019), a 2/3 majority of voting members was not reached. Primary objection to the changes focused on remaining uncertainties in some relationships and the potential confusion that may be created if the proposed changes require further revision in the near future. At issue are 1) the conflicting support among data sets for a sister-taxa relationship between *L. albirostris* and *L. acutus* (which

would obviate the need for *Leucopleurus*), and 2) evidence that *australis* and *cruciger* should be included in *Cephalorhynchus* (which would necessitate a new genus for *obliquidens* and *obscurus*, as *australis* is the type species for the genus *Sagmatias*). Therefore, the taxonomy list retains all species in *Lagenorhynchus*, recognizing that the current taxonomy does not reflect the evolutionary distinctiveness of *L. albirostris* and *L. acutus*. Next steps in unraveling the taxonomy of these dolphins will need to involve robust datasets that include all *Lagenorhynchus* and *Cephalorhynchus* species. Harlin-Cognato (2010) recognized *L. obscurus posidonia* (Peru/Chile).

It has been noted repeatedly, most recently by Perrin et al. (2013), that the delphinine genera *Stenella* and *Tursiops* are paraphyletic and that at present there is no molecular or morphological basis for satisfactory resolution of phylogenetic relationships in the subfamily. A possible solution would be to return all the species in *Tursiops*, *Sousa*, and *Stenella* to *Delphinus*, the genus in which they were first described, and place *Lagenodelphis hosei* there as well. However, considering that this would obscure the clear close relationship of the present *Sousa* and *Tursiops* species, the status quo is maintained here provisionally, pending the outcome of more definitive morphological and molecular studies. Eventually a more natural classification will emerge.

Perrin et al. (1999) established the subspecies *Stenella longirostris roseiventris*.

The Irrawaddy dolphin was split into *O. brevirostris* and *O. heinsohni*, the Australian snubfin dolphin (Beasley et al. 2005).

Wang et al. (2008) and Jefferson and Wang (2011) established *Neophocaena asiaorientalis* as a full species distinct from the *N. phocaenoides*, with two subspecies. Zhou et al. (2018) examined genomic DNA sequence data and concluded that there was significant genetic differentiation between the two subspecies of *N. asiaorientalis*, including evidence for genetic adaptation to the freshwater environment of the Yangtze River. The authors proposed the two subspecies of *N. asiaorientalis* be elevated to species-level, the Yangtze finless porpoise, *N. asiaorientalis* Pilleri and Gühr 1927, and the East Asian finless porpoise, *N. sunameri* Pilleri and Gühr 1972. However, there were no samples in this study from near the mouth of the Yangtze River where the marine and river subspecies could possibly be sympatric and where genetic interchange could occur. While the results of the study are consistent with recently diverged taxa, the lack of samples in the potential area of sympatry make it difficult to rule out an isolation by distance effect.

Viaud-Martinez et al. (2007) concluded based on morphological and molecular genetic evidence that *Phocoena phocoena relicta* is a distinct subspecies.

We list the baiji *Lipotes vexillifer* as “possibly extinct” in conformance with the IUCN Red List, although extinction seems a certainty.

Manatees and Dugongs

In the Sirenia, subspecies of the dugong are not currently recognized (Domning, 1996; see <http://sirendom.org/biblio> for update to 1996 publication). However, no in-depth study has been undertaken to address the issue of subspecies.

## References

- Agnarsson, I. and L. J. May-Collado. 2008. The phylogeny of Cetartiodactyla: The importance of dense taxon sampling, missing data, and the remarkable promise of cytochrome *b* to provide reliable species-level phylogenies. *Molecular Phylogenetics and Evolution* 48:964–985.
- Archer F.I., R. L. Brownell Jr, B. L. Hancock-Hanser, P. A. Morin, K. M. Robertson, K. K. Sherman, J. Calambokidis, J. Urbán, P. E. Rosel, S. A. Mizroch, S. Panigada, B. L. Taylor. 2019. Subspecies of fin whales (*Balaenoptera physalus* Linnaeus, 1758): Taxonomic implications of genetics. *Journal of Mammalogy* 100:1653–1670.
- Baker, R. J., L. C. Bradley, R. D. Bradley, J. W. Dragoo, M. D., R. S. Hoffman, C. A. Jones, F. Reid, D. W. Rice and C. Jones. 2003. Revised checklist of North American mammals north of Mexico, 2003. *Museum of Texas Tech University Occasional Papers* 229:1–24.
- Banguera-Hinestroza, E., M. Cárdenas, M. Ruiz-García, M. Marmontel, E. Gaitán, R. Vázquez and F. García-Vallejo. 2002. Molecular identification of evolutionarily significant units in the Amazon river dolphin *Inia* sp. (Cetacea: Iniidae). *Journal of Heredity* 93:312–322.
- Banguera-Hinestroza, E., A. Hayano, E. Crespo and A. R. Hoelzel. 2014. Delphinid systematics and biogeography with a focus on the current genus *Lagenorhynchus*: multiple pathways for antitropical and trans-oceanic radiation. *Molecular Phylogenetics and Evolution*, 80:217–230.
- Banks, R. C. and R. L. Brownell. 1969. Taxonomy of the common dolphins of the eastern Pacific Ocean. *Journal of Mammalogy* 50:262–271.
- Beasley, I., K. M. Robertson and P. Arnold. 2005. Description of a new dolphin, the Australian snubfin dolphin *Orcaella heinsohni* sp. n. (Cetacea: Delphinidae). *Marine Mammal Science* 21:365–400.
- Berta, A. 1994. New specimens of the pinnipediform *Pteronarctos* from the Miocene of Oregon. *Smithsonian Contributions to Paleobiology* no. 78:1–33.
- Berta, A. and M. Churchill. 2012. Pinniped taxonomy: Review of currently recognized species and subspecies, and evidence used for their description. *Mammal Review* 42:207–234.
- Berta, A. and A. R. Wyss. 1994. Pinniped phylogeny. Pages 33–56 in A. Berta and T. A. Deméré (eds). *Contributions in Marine Mammal Paleontology Honoring Frank C. Whitmore, Jr.* Proceedings of the San Diego Society of Natural History. San Diego, California, USA.

Branch, T. A., E. M. N. Abubaker S. Mkango S and D. S. Butterworth. 2007. Separating southern blue whale subspecies based on length frequencies of sexually mature females. *Marine Mammal Science* 23:803–833.

Braulik, G.T., F.I. Archer, U. Khan, M. Imran, R.K. Sinha, T.A. Jefferson, C. Donovan, C. and J.A. Grave. 2021. Taxonomic revision of the South Asian River dolphins (*Platanista*): Indus and Ganges River dolphins are separate species. *Marine Mammal Science* 37:1022–1059.

Braulik, G. T., R. Barnett, V. Odon, V. Islas-Villanueva, A. R. Hoelzel and J. A. Graves. 2014. One species or two? Vicariance, lineage divergence and low mtDNA diversity in geographically isolated populations of South Asia river dolphin. *Journal of Mammal Evolution* 22:111–120.

Brunner, S. 2004. Fur seals and sea lions (Otariidae): Identification of species and taxonomic review. *Systematics and Biodiversity* 1:339–439.

Buchan, S. J., R. Huckle-Gaete, K. M. Stafford and C. W. Clark. 2018. Occasional acoustic presence of Antarctic blue whales on a feeding ground in southern Chile. *Marine Mammal Science* 34:220–228.

Cañizales, I. 2020. Morphology of the skull of *Inia geoffrensis humboldtiana* Pilleri & Gühr, 1977 (Cetacea:Iniidae): A morphometric and taxonomic analysis. *Graellsia*, 76(2): e115.

Carroll, E.L., M. R. McGowen, M. L. McCarthy, F. G. Marx, N. Aguilar, M. L. Dalebout, S. Dreyer, O. E. Gaggiotti, S. S. Hansen, A. van Helden, A. B. Onoufriou, R. W. Baird, C. S. Baker, S. Berrow, D. Cholewiak, D. Claridge, R. Constantine, N. J. Davison, C. Eira, R. E. Fordyce, J. Gatesy, G. J. Greg Hofmeyr, V. Martín, J. G. Mead, A. A. Mignucci-Giannoni, P. A. Morin, C. Reyes, E. Rogan, M. Rosso, M. A. Silva, M. S. Springer, D. Steel and M. Tange Olsen. 2021. Speciation in the deep: genomics and morphology reveal a new species of beaked whale *Mesoplodon eueu*. *Proceedings of the Royal Society B*, 288:20211213.

Charlton-Robb, K., L. Gershwin, R. Thompson, J. Austin, K. Owen and S. McKechnie. 2011. A new dolphin species, the Burrnanan dolphin *Tursiops australis* sp. Nov., endemic to southern Australian waters. *PLoS ONE* 6(0) e24047.

Cipriano, F. 1997. Antitropical distributions and speciation in dolphins of the genus *Lagenorhynchus*: A preliminary analysis. Pages 305–316 in A. E. Dizon, S. J. Chivers and W. F. Perrin (eds). *Molecular genetics of marine mammals*. Society for Marine Mammalogy Special Publication 3.

Clarke, R. 2004. Pygmy fin whales. *Marine Mammal Science* 20:329–334.

Costa, A. P. B., P. E. Rosel, F. G. Daura-Jorge and P. C. Simões-Lopes. 2016. Offshore and coastal common bottlenose dolphins of the western South Atlantic face-to-face: What the skull and the spine can tell us. *Marine Mammal Science* 32:1433–1457.

Costa, A. P. B., P. F. Fruct, E. R. Secchi, F. G. Daura-Jorge, P. C. Simões-Lopes, J. C. Di Tullio and P. E. Rosel. 2021. Ecological divergence and speciation in common bottlenose dolphins in the western South Atlantic. *Journal of Evolutionary Biology* 34:16–32.

Costa, A. P., W. Mcfee, L. A. Wilcox, F. I Archer and P. E. Rosel. 2022. The common bottlenose dolphin (*Tursiops truncatus*) ecotypes of the western North Atlantic revisited: an integrative taxonomic investigation supports the presence of distinct species. *Zoological Journal of the Linnean Society*, 196(4), 1608-1636.

Costa, A. P. B., F. I. Archer, P. E. Rosel, and W. F. Perrin. 2023. *Tursiops truncatus nuuanu*, a new subspecies of the common bottlenose dolphin from the eastern tropical Pacific. *Journal of Mammalian Evolution* 30:213–229

Cunha, H. A., R. Loizaga de Castro, E. R. Secchi, E. A. Crespo, J. Lailson-Brito, A. F. Azevedo, C. Lazoski and A. M. Solé-Cava. 2015. Molecular and morphological differentiation of common dolphins (*Delphinus* sp.) in the southwestern Atlantic: Testing the two species hypothesis in sympatry. *PLoS ONE* 10(11) e0140251.

Dalebout, M. L., J. G. Mead, C. S. Baker, A. N. Baker and A. L. van Helden. 2002. A new species of beaked whale *Mesoplodon perrini* sp. n. (Cetacea: Ziphiidae) discovered through phylogenetic analyses of mitochondrial DNA sequences. *Marine Mammal Science* 18:577–608.

Dalebout, M. L., S. C. Baker, D. Steel, K. Thompson, K. M. Robertson, S. J. Chivers, W. F. Perrin, M. Goonatilake, R. C. Anderson, J. G. Mead, C. W. Potter, L. Thompson, D. Jupiter and T. K. Yamada. 2014. Resurrection of *Mesoplodon hotaula* Deraniyagala 1963: A new species of beaked whale in the tropical Indo-Pacific. *Marine Mammal Science* 30:1081–1108.

Derocher, A. E. and I. Stirling. 1998. Geographic variation in growth of polar bears (*Ursus maritimus*). *Journal of Zoology* 245:65–72.

Domning, D. 1996. Bibliography and index of the Sirenia and Desmostylia. *Smithsonian Contributions to Paleobiology* 80:1–611.

Emin-Lima, R., F. A. Machado, S. Siciliano, W. Gravena, E. Aliaga-Rossel, J. de Sousa e Silva, Jr., E. Hingst-Zaher and L. R. de Oliveira, L. R. 2022. Morphological disparity in the skull of Amazon River dolphins of the genus *Inia* (Cetacea, Iniidae) is inconsistent with a single taxon. *Journal of Mammalogy*, 103(6), 1278-1289.

Fordyce, R. E. and F. G. Marx. 2012. The pygmy right whale *Caperea marginata*: Last of the cetotheres. *Proceedings of the Royal Society B* 280:20122645.

Frère, C. H., P. Hale, L. Porter, V. G. Cockcroft and M. L. Dalebout. 2008. Phylogenetic analysis of mtDNA sequences suggests revision of humpback dolphin (*Sousa* spp.) taxonomy is needed. *Marine and Freshwater Research* 59:259–268.

Frère, C. H., J. Seddon, C. Palmer, L. Porter and G. J. Parra. 2011. Multiple lines of evidence for an Australasian geographic boundary in the Indo-Pacific humpback dolphin (*Sousa chinensis*): Population or species divergence? *Conservation Genetics* 12:1633–1638.

Fulton, T. L. and C. Strobeck. 2010. Multiple fossil calibrations, nuclear loci and mitochondrial genomes provide new insight into biogeography and divergence timing for true seals (Phocidae, Pinnipedia). *Journal of Biogeography* 37:814–829.

Gaines, C. A., M. P. Hare, S. E. Beck and H. C. Rosenbaum. 2005. Nuclear markers confirm taxonomic status and relationships among highly endangered and closely related right whale species. *Proceedings of the Royal Society B* 272:533–542.

Gardner, A. L. and C. B. Robbins. 1999. Case 3058. *Arctocephalus* F. Cuvier, 1826 and *Callorhinus* Gray, 1859 (Mammalia, Pinnipedia): proposed conservation by the designation of *Phoca pusilla* Schreber, [1775] as the type species of *Arctocephalus*; and *Otaria* Péron, 1816 and *Eumetopias* Gill, 1866: proposed conservation by the designation of *Phoca leonina* Molina, 1782 as the type species of *Otaria*. *Bulletin of Zoological Nomenclature* 56:136–141.

Geisler, J. H. and M. D. Uhen. 2005. Phylogenetic relationships of extinct cetartiodactyls: Results of simultaneous analyses of molecular, morphological, and stratigraphic data. *Journal of Mammalian Evolution* 12:145–160.

Graur D., and D. G. Higgins. 1994. Molecular evidence for the inclusion of cetaceans within the order Artiodactyla. *Molecular Biology and Evolution* 11:357–364.

Gravena, W., I.P. Farias, M.N.F. da Silva, V.M.F. da Silva and T. Hrbek. 2014. Looking into the past and the future: were the Madeira River rapids a geographical barrier to the boto (Cetacea: Iniidae)? *Conservation Genetics* 15:619–629.

Harlin-Cognato, A. D. 2010. The dusky dolphin's place in the delphinid family tree. Pages 1–20 in B. Würsig and M. Würsig (eds). *The dusky dolphin. Master acrobat off different shores*. Academic Press, Amsterdam.

van Helden, A. L., A. N. Baker, M. L. Dalebout, J. C. Reyes, K. Van Waerebeek, and C. S. Baker. 2002. Resurrection of *Mesoplodon traversii* (Gray, 1874), senior synonym of *M. bahamondi* Reyes, Van Waerebeek, Cárdenas and Yáñez, 1995 (Cetacea: Ziphiidae). *Marine Mammal Science* 18:609–621.

Heyning, J. E. and W. F. Perrin. 1994. Evidence for two species of common dolphins (genus *Delphinus*) from the eastern North Pacific. *Contributions to Science of the Natural History Museum of Los Angeles City* 442:1–35.

Hershkovitz, P. 1966. *Catalog of living whales*. United States National Museum Bulletin 246, 259pp.



- Higdon, J. W., O. R. P. Bininda-Emonds, R. M. Beck and S. H. Ferguson. 2007. Phylogeny and divergence of the pinnipeds (Carnivora, Mammalia) assessed using a multigene dataset. *BMC Evolutionary Biology* 7:216.
- Hohl, L. S., F. L. Sicuro, J. C. Wickert, I. B. Moreno, O. Rocha-Barbosa and A. S. Barreto. 2020. Skull morphology of bottlenose dolphins from different ocean populations with emphasis on South America. *Journal of Morphology* 281:564–577
- Hollatz, C., S. Torres Vilaça, R. A. F. Redondo, M. Marmontel, C. S. Baker and F. R. Santos. 2011. The Amazon River system as an ecological barrier driving genetic differentiation of the pink dolphin (*Inia geoffrensis*). *Biological Journal of the Linnaen Society* 102:812–827.
- Hrbek, T., V. M. F. da Silva, N. Dutra, W. Gravena, A. R. Martin and I. P. Farias. 2014. A new species of river dolphin from Brazil or: How little do we know our biodiversity. *PLoS ONE* 9(1) e83623.
- International Committee on Zoological Nomenclature [ICZN]. 1999. International Code of Zoological Nomenclature. Fourth Edition. International Trust for Zoological Nomenclature c/o The Natural History Museum – Cromwell Road, London, United Kingdom.
- International Whaling Commission. 2001. Report of the Scientific Committee. *Journal of Cetacean Research and Management* 3, Supplement:1–75.
- Jackson, J. A., D. J. Steel, P. Beerli, B. C. Congdon, C. Olavarria, M. S. Leslie, C. Pomilla, H. Rosenbaum and C. S. Baker. 2014. Global diversity and oceanic divergence of humpback whales (*Megaptera novaeangliae*). *Proceedings of the Royal Society B* 281, 20133222:1–10.
- Jedensjö M., C. M. Kemper, M. Milella, E. P. Willems, M. Krützen. 2020. Taxonomy and distribution of bottlenose dolphins in Australian waters: an osteological clarification. *Canadian Journal of Zoology* 98:461–479
- Jefferson, T.A. and H. C. Rosenbaum. 2014. Taxonomic revision of the humpback dolphins (*Sousa* spp.), and description of a new species from Australia. *Marine Mammal Science* 30:1494–1541.
- Jefferson, T. A. and J. Y. Wang. 2011. Revision of the taxonomy of finless porpoises (genus *Neophocaena*): The existence of two species. *Journal of Marine Animals and Their Ecology* 4:3–16.
- Kato, H. and W. F. Perrin. 2009. Bryde's whales *Balaenoptera edeni/brydei*. Pages 158–163 in W. F. Perrin, B. Würsig and J. G. M. Thewissen (eds) *Encyclopedia of Marine Mammals*. Academic Press, Amsterdam.
- Kershaw, F., M. S. Leslie, T. Collins, R. M. Mansur, B. D. Smith, G. Minton, R. Baldwin, R. G. LeDuc, R. C. Anderson, R. L. Brownell Jr and H. C. Rosenbaum. 2013. Population

differentiation of 2 forms of Bryde's whales in the Indian and Pacific Oceans. *Journal of Heredity* 6:755–764.

Khalif, N. 2020. The Chilean blue whale (*Balaenoptera musculus chilensis* Khalaf, 2020): A new subspecies from Chile. *Gazelle: The Palestinian Biological Bulletin* 38: 40–46.

Khalif, N. 2021. The Arabian Sea blue whale (*Balaenoptera musculus arabica* Khalaf, 2021): A new subspecies from the Arabian Sea and western Indian Ocean. *Gazelle: The Palestinian Biological Bulletin* 39:17–46.

Kinze, C. C. 2018. A case for *Tursiops tursio* (Gunnerus, 1768). *Lutra* 61:189–196.

Kitamura, S., T. Matsuishi, T. K. Yamada, Y. Tajima, H. Ishikawa, S. Tanabe, H. Nakagawa, Y. Uni and S. Abe. 2013. Two genetically distinct stocks in Baird's beaked whale (Cetacea: Ziphiidae). *Marine Mammal Science* 29:755–766.

Koretsky, I. A., L. G. Barnes and S. J. Rahmat. 2016. Re-evaluation of morphological characters questions current views of pinniped origins. *Vestnik Zoologii* 50:327–354.

Larivière, S. 1998. *Lontra felina*. *Mammalian Species* 575:1–5.

LeDuc, R. G., W. F. Perrin and A. E. Dizon. 1999. Phylogenetic relationships among the delphinid cetaceans based on full cytochrome b sequences. *Marine Mammal Science* 15:619–648.

LeDuc, R. G., F. I. Archer, A. R. Lang, K. K. Martien, B. Hancock-Hanser, J. O. Torres-Florez, R. Hucke-Gaete, H. C. Rosenbaum, K. Van Waerebeek, R. L. Brownell Jr and B. L. Taylor. 2017. Genetic variation in blue whales in the eastern Pacific: implication for taxonomy and use of common wintering grounds. *Molecular Ecology* 26:740–751.

LeDuc, R. G., A. E. Dizon, M., Goto, M., L. A. Pastene, H. Kato, S. Nishiwaki and R. L. Brownell 2007. Patterns of genetic variation in southern hemisphere blue whales, and the use of assignment test to detect mixing on the feeding grounds. *Journal of Cetacean Research and Management* 9:73–80.

Leslie, M. S., C. M. Perkins-Taylor, J. W. Durban, M. J. Moore, C. A. Miller, P. Chanarat, P. Bahamonde, G. Chiang and A. Apprill. 2020. Body size data collected non-invasively from drone images indicate a morphologically distinct Chilean blue whale (*Balaenoptera musculus*) taxon. *Endangered Species Research* 43:291–304.

Lindqvist, C., L. Bachmann, L. W. Andersen, E. W. Born, U. Arnason, K. M. Kovacs, C. Lydersen, A. V. Abramov and Ø. Wiig. 2009. The Laptev Sea walrus *Odobenus rosmarus laptevi*: An enigma revisited. *Zoologica Scripta* 38:113–127.

- Luksenburg, J. A., A. Henriquez and G. Sangster. 2015. Molecular and morphological evidence for the subspecific identity of Bryde's whales in the southern Caribbean. *Marine Mammal Science* 31:1568–1579.
- Marx, F. G. and R. E. Fordyce. 2015. Baleen whale boom and bust: A synthesis of mysticete phylogeny, diversity and disparity. *Royal Society Open Science* 2: 140434.
- Marx, F.G. and R. E. Fordyce. 2016. A link no longer missing: New evidence for the cetotheriid affinities of *Caperea*. *PLoS ONE* 11(10) e0164059.
- McDonald, M. A., S. L. Mesnick and J. A. Hildebrand. 2006. Biogeographic characterization of blue whale song worldwide: using song to identify populations. *Journal of Cetacean Research and Management* 8:55–65.
- McGowen, M. R. 2011. Toward the resolution of an explosive radiation—A multilocus phylogeny of oceanic dolphins (Delphinidae). *Molecular Phylogenetics and Evolution* 60:345–357.
- Mendez, M., T. A. Jefferson, S.-O. Kolokotronis, M. Krützen, G. J. Parra, T. Collins, G. Minton, R. Baldwin, P. Berggren, A. Särnblad, O. A. Amir, V. M. Peddemors, L. Karczmarski, A. Guissamulo, B. Smith, D. Sutaria, G. Amato and H. C. Rosenbaum. 2013. Integrating multiple lines of evidence to better understand the evolutionary divergence of humpback dolphins along their entire distribution range: A new dolphin species in Australian waters. *Molecular Ecology* 22:5936–5948.
- Montgelard C., F. M. Catzeflis and E. Douzery. 1997. Phylogenetic relationships of artiodactyls and cetaceans as deduced from the comparison of cytochrome I and 12S rRNA mitochondrial sequences. *Molecular Biology and Evolution* 14:550-559.
- Morin, P. A., C. S. Baker, R. S. Brewer, A. M. Burdin, M. L. Dalebout, J. P. Dines, I. Fedutin, O. Filatova, E. Hoyt, J. L. Jung and M. Lauf. 2017. Genetic structure of the beaked whale genus *Berardius* in the North Pacific, with genetic evidence for a new species. *Marine Mammal Science* 33:96–111.
- Moura, A. E., K. Shreves, M. Pilot, K. R. Andrews, D. M. Moore, T. Kishida, L. Möller, A. Natoli, S. Gaspari, M. McGowen, I. Chen, H. Gray, M. Gore, R. M. Culloch, M. S. Kiani, M. S. Willson, A. Bulushi, T. Collins, R. Baldwin, A. Willson, G. Minton, L. Ponnampalam and A. R. Hoelzel. 2020. Phylogenomics of the genus *Tursiops* and closely related Delphininae reveals extensive reticulation among lineages and provides inference about eco-evolutionary drivers. *Molecular Phylogenetics and Evolution* 146:106756.
- Natoli, A., A. Cañadas, V. M. Peddemors, A. Aguilar, C. Vaquero, P. Fernandez-Piqueras and A. R. Hoelzel. 2006. Phylogeography and alpha taxonomy of the common dolphin (*Delphinus* sp.). *Journal of Evolutionary Biology* 19:943–954.

- Natoli, A., A. Cañadas, C. Vaquero, E. Politi, P. Fernandez-Navarro and A. R. Hoelzel. 2008. Conservation genetics of the short-beaked common dolphin (*Delphinus delphis*) in the Mediterranean Sea and the eastern North Atlantic. *Conservation Genetics* 9:1479–1487.
- Nyakatura, K. and O. R.P. Birinda-Emonds. 2012. Updating the evolutionary history of Carnivora (Mammalia): A new species-level supertree complete with divergence time estimates. *BMC Biology* 10:1–31.
- Oliveira, L. R. and R. L. Brownell, Jr. 2014. Taxonomic status of two subspecies of South American fur seals: *Arctocephalus australis australis* vs. *A. a. gracilis*. *Marine Mammal Science* 30:1258–1263.
- Oliveira, L. R. D., L. D. Fraga, P. H. Ott, S. Siciliano, F. Lopes, R. Almeida, J. C. Wickert, L. Milmann, D. Danilewicz, N. R. Emin-Lima and A. C. Meirelles. 2019. Population structure, phylogeography, and genetic diversity of the common bottlenose dolphin in the tropical and subtropical southwestern Atlantic Ocean. *Journal of Mammalogy*, 100:564–577.
- Olsen, M. T., A. Galatius, V. Biard, K. Gregersen and C. C. Kinze. 2016. The forgotten type specimen of the grey seal [*Halichoerus grypus* (Fabricius, 1791)] from the island of Amager, Denmark. *Journal of the Linnean Society* 2016:1–8.
- Orr, H. A. and J. A. Coyne. 2004. *Speciation*. Sinauer Associates, Sunderland, Massachusetts.
- Pérez-Alvarez, M., S. Kraft, N. I. Segovia, C. Olavarria, S. Nigenda-Morales, R. J. Urbán, L. Vilorio-Gómora, F. Archer, R. Moraga, M. Sepúlveda, and M. Santos-Carvalho. 2021. Contrasting phylogeographic patterns among Northern and Southern Hemisphere fin whale populations with new data from the Southern Pacific. *Frontiers in Marine Science* 8:630233.
- Perrin, W. F. 2009. Species. Pages 1084–1087 in W. F. Perrin, B. Würsig and J. G. M. Theissen (eds) *Encyclopedia of Marine Mammals*. Academic Press, Amsterdam.
- Perrin, W. F., M. L. L. Dolar and D. Robineau. 1999. Spinner dolphins (*Stenella longirostris*) of the western Pacific and Southeast Asia: Pelagic and shallow-water forms. *Marine Mammal Science* 15:1029–1053.
- Perrin, W. F., J. L. Thieleking, W. A. Walker, F. I. Archer and K. M. Robertson. 2011. Common bottlenose dolphins (*Tursiops truncatus*) in California waters: Cranial differentiation of coastal and offshore ecotypes. *Marine Mammal Science* 27:769–792.
- Perrin, W. F., P. E. Rosel and F. Cipriano. 2013. How to contend with paraphyly in the taxonomy of the delphinine cetaceans. *Marine Mammal Science* 29:567–588.
- Price, S. A., O. R. P. Bininda-Emonds and J. L. Gittleman. 2005. A complete phylogeny of the whales, dolphins and even-toed hoofed mammals (Cetartiodactyla). *Biological Review* 80:445–473.

- Prothero, D. R., D. Domning, R. E. Fordyce, S. Foss, C. Janis, S. Lucas, K. L. Marriott, G. Metais, D. Naish, K. Padian and G. Rössner. 2022. On the unnecessary and misleading taxon “Cetartiodactyla”. *Journal of Mammalian Evolution* 29:93-97.
- de Queiroz, K. 2007. Species concepts and species delineation. *Systematic Biology* 56:879–886.
- Reeves, R. R., W. F. Perrin, B. L. Taylor, C. S. Baker and S. L. Mesnick (eds) 2004. Report of the Workshop on Shortcomings of Cetacean Taxonomy in Relation to Needs of Conservation and Management, April 30–May 2, 2004 La Jolla, California. NOAA Technical Memorandum NOAA-TM-NMFS-SWFSC-363:1–94.
- Rice, D. W. 1998. Marine mammals of the world. Systematics and distribution. Society for Marine Mammalogy Special Publication 4:1–231.
- Rosel, P. E. and L. A. Wilcox. 2014. Genetic evidence reveals a unique lineage of Bryde’s whales in the northern Gulf of Mexico. *Endangered Species Research* 23:19–34.
- Rosel, P. E., L. A. Wilcox, T. K. Yamada and K. D. Mullin. 2021. A new species of baleen whale (*Balaenoptera*) from the Gulf of Mexico, with a review of its geographic distribution. *Marine Mammal Science* 37:577–610.
- Rosel, P. E., A. E. Dizon and J. E. Heyning. 1994. Genetic analysis of sympatric morphotypes of common dolphins (genus *Delphinus*). *Marine Biology* 119:150–167.
- Rosenbaum, H., R. L. Brownell, Jr., M. W. Brown, C. Schaeff, Y. Portway, B. N. White, S. Malik, L. A. Pastene, N. J. Patenaude, C. S. Baker, M. Goto, P. B. Best, P. J. Clapham, P. Hamilton, R. Payne, V. Rowntree, C. T. Tynan, J. L. Bannister, and R. DeSalle. 2000. World-wide genetic differentiation of *Eubalaena*: Questioning the number of right whale species. *Molecular Ecology* 9:1793–1802.
- Ruiz-García, M., E. Banguera and H. Cárdenas. 2006. Morphological analysis of three *Inia* (Cetacea: Iniidae) populations from Colombia and Bolivia. *Acta Theriologica* 51:411–426.
- Ruiz-García, M., S. Caballero, M. Martínez-Agüero and J. M. Shostell. 2008. Molecular differentiation among *Inia geoffrensis* and *Inia boliviensis* (Iniidae, Cetacea) by means of nuclear intron sequences. Pages 1–25 in V. T. Koven. ed., *Population Genetics Research Progress* Chapter 6.
- Sasaki, T., M. Nikaido, S. Wada, T. K. Yamada, Y. Cao, M. Hasegawa and N. Okada. 2006. *Balaenoptera omurai* is a newly discovered baleen whale that represents an ancient evolutionary lineage. *Molecular Phylogenetics and Evolution* 41:40–52.
- Scheel, D.-M., G. J. Slater, S.-O. Kolokontronis, C. W. Potter, D. S. Rotstein, K. Tsangaras, A. D. Greenwood and K. M. Helgen. 2014. Biogeography and taxonomy of extinct and endangered monk seals illuminated by ancient DNA and skull morphology. *ZooKeys* 409:1–33.

- Siciliano, S., V. H. Valiati, R. Emin-Lima, A. F. Costa, J. Sartor, T. Dorneles and L. R. De Oliveira. 2016. New genetic data extend the range of river dolphins *Inia* in the Amazon Delta. *Hydrobiologia* 777:255–269.
- da Silva, V. M. F. 1994. Aspects of the biology of the Amazonian dolphin genera *Inia* and *Sotalia fluviatilis*. Ph.D. thesis, University of Cambridge, Cambridge, U.K. 328 pp.
- Spaulding, M., M. A. O’Leary and J. Gatesy. 2009. Relationships of Cetacea (Artiodactyla) among mammals: Increased taxon sampling alters interpretations of key fossils and character evolution. *PLoS ONE* 4(9) e7062.
- Taylor, B. L., F. I. Archer, K. K. Martien, P. E. Rosel, B. L. Hancock-Hanser, A. R. Lang, M. S. Leslie, S. L. Mesnick, P. A. Morin, V. L. Pease, W. F. Perrin, K. M. Robertson, K. M. Parsons, A. Viricel, N. L. Vollmer, R. R. Reeves, F. Cipriano, M. Krützen and C. S. Baker. 2017a. Guidelines and quantitative standards to improve consistency in cetacean subspecies and species delimitation relying on molecular genetic data. *Marine Mammal Science* 33:132–155.
- Taylor B. L., W. F. Perrin, R. R. Reeves, P. E. Rosel, J. Y. Wang, F. Cipriano, C. S. Baker and R. L. Brownell Jr. 2017b. Why we should develop guidelines and quantitative standards for using molecular genetic data to delimit subspecies for data-poor organisms like cetaceans. *Marine Mammal Science* 33:12–26.
- Thompson, K., C. S. Baker, A. van Helden, S. Paatel, C. Millar and R. Constantine. 2012. The world’s rarest whale. *Current Biology* 22:R905–R906.
- Torres-Florez, J. P., R. Hucke-Gaete, R. LeDuc, A. Lang, B. Taylor, L. E. Pimper, L. Bedriñana-Romano, H. C. Rosenbaum, and C. C. Figueroa. 2014. Blue whale population structure along the eastern South Pacific Ocean: evidence of more than one population. *Molecular Ecology* 23:5998–601.
- Viaud-Martinez, M. Martinez Vergara, P. E. Gol’din, V. Ridoux, A. A. Öztürk, B. Öztürk, P. E. Rosel, A. Frantzis, A. Komnenou and A. J. Bohanak. 2007. Morphological and genetic differentiation of the Black Sea harbour porpoise *Phocoena phocoena*. *Marine Ecology Progress Series* 338:281–294.
- Viaud-Martinez, K. A., R. L. Brownell, Jr., A. Komnenou and A. J. Bohanak. 2008. Genetic isolation and morphological divergence of Black Sea bottlenose dolphins. *Biological Conservation* 141:1600–1611.
- Vollmer, N. L., E. Ashe, R. L. Brownell Jr, F. Cipriano, J. G. Mead, R. R. Reeves, M. S. Soldevilla and R. Williams, R. 2019. Taxonomic revision of the dolphin genus *Lagenorhynchus*. *Marine Mammal Science*, 35: 957–1057.
- Wada, S., M. Oishi and T. K. Yamada. 2003. A newly discovered species of living baleen whale. *Nature* 426:278–281.

Wang, J. Y., T. R. Frasier, S. C. Yang and B. N. White. 2008. Detecting recent speciation events: The case of the finless porpoise (genus *Neophocaena*). *Heredity* 101:145–155.

Wang, J. Y., S. C. Yang and S. K. Hung. 2015. Diagnosability and description of a new subspecies of Indo-Pacific humpback dolphin, *Sousa chinensis*, (Osbeck, 1765) from the Taiwan Strait. *Zoological Studies* 54:36.

Wang, J.Y., A.P.B. Costa and T.A. Jefferson. 2021. The correct name of Lahille's bottlenose dolphin, *Tursiops truncatus gephyreus* Lahille, 1908. *Marine Mammal Science* 37: 696–701.

Webber, M. S. 2014. Family Otariidae (eared seals). Pages 84–101 in D. E. Wilson and R. A. Mittermeier (eds). *Handbook of the Mammals of the World. 4. Sea Mammals*. Lynx Ediciones, Barcelona.

Wickert, J. C., S. M. Von Eye, L. R. Oliveira and I. B. Moreno. 2016. Revalidation of *Tursiops gephyreus* Lahille, 1908 (Cetartiodactyla: Delphinidae) from the southwestern Atlantic Ocean. *Journal of Mammalogy* 97:1728–1737.

Wyss, A. R. and J. Flynn. 1993. A phylogenetic analysis and definition of the Carnivora. Pages 32–52 in F. S. Szalay, M. J. Novacek, and M. C. McKenna (eds). *Mammal Phylogeny: Placentals*. Springer-Verlag, New York.

Yamada, T. K., S. Kitamura, S. Abe, Y. Tajima, A. Matsuda, J. G. Mead and T.F. Matsuishi. 2019. Description of a new species of beaked whale (*Berardius*) found in the North Pacific. *Scientific Reports* 9:1–14.

Last updated June 2023 by members of the [Committee on Taxonomy](#):

- Patricia E. Rosel (Chair)
- Frederick I. Archer
- C. Scott Baker
- Daryl J. Boness
- Robert L. Brownell, Jr.
- Morgan Churchill
- Ana P. Costa
- Daryl P. Domning
- Thomas A. Jefferson
- Carl Kinze
- Michael McGowen
- Larissa R. Oliveira
- John Y. Wang
- Tadasu K. Yamada